

Author Index

- Aman, J. *see* Lodefalk, M.
- Abbas, J. *see* Forouhi, N.G.
- Abdella, N.A. *see* Mojiminiyi, O.A.
- Acerini, C.L. Weight matters for children. A complete guide for weight, eating and fitness, 1275
- Adaikalakoteswari, A., Balasubramanyam, M., Rema, M. & Mohan, V. Differential gene expression of NADPH oxidase (p22^{phox}) and hemoxygenase-1 in patients with Type 2 diabetes and microangiopathy, 666
- Adamopoulos, I. *see* Angelopoulos, N.G.
- Adang, E. *see* Dijkstra, R.F.
- Agarwal, M.M., Dhatt, G.S. & Punnose, J. Gestational diabetes: utility of fasting plasma glucose as a screening test depends on the diagnostic criteria, 1319
- Agius, E., Attard, G., Shakespeare, L., Clark, P., Vidya, M.-A., Hattersley, A.T. & Fava, S. Familial factors in diabetic nephropathy: an offspring study, 331
- Aguilar-Salinas, C.A. *see* Navalón-García, K.
- Ahmad, N. *see* Lawton, J.
- Ahmed, A.B. *see* Khan, A.S.A.
- Ahmed, A.T., Karter, A.J. & Liu, J. Alcohol consumption is inversely associated with adherence to diabetes self-care behaviours, 795
- Ahn, C.W. *see* Cho, N.H.
- Ahn, J.H., Kim, T.Y., Kim, Y.-J., Han, M.W., Yoon, T.H. & Chung, J.W. Lipo-prostaglandin E₁ in combination with steroid therapy is effective for treatment of sudden sensorineural hearing loss in Korean patients with Type 2 diabetes, 1339
- Ahn, T.Y. *see* Cho, N.H.
- Ajmani, A. *see* Vikram, N.K.
- Akram, K., Pedersen-Bjergaard, U., Carstensen, B., Borch-Johnsen, K. & Thorsteinsson, B. Frequency and risk factors of severe hypoglycaemia in insulin-treated Type 2 diabetes: a cross-sectional survey, 750
- Aksoy, M. *see* Ertugrul, M.B.
- Alberti, H. *see* Harzallah, F.
- Alberti, H. Diabetes in the Arab World, 107
- Alberti, K.G.M.M. *see* Whincup, P.
- Alberti, K.G.M.M., Zimmet, P. & Shaw, J. Metabolic syndrome—a new world-wide definition. A Consensus Statement from the International Diabetes Federation, 469
- Alberti, K.G.M.M., Zimmet, P. & Shaw, J. Reply, 1385
- Albrechtsen, A. *see* Wegner, L.
- Aldhahi, W.A. *see* Mojiminiyi, O.A.
- Alem, M. *see* Nazir, F.S.
- Ali, S., Stone, M.A., Peters, J.L., Davies, M.J. & Khunti, K. The prevalence of co-morbid depression in adults with Type 2 diabetes: a systematic review and meta-analysis, 1165
- Allard, S.E. *see* Browne, D.L.
- Allard, S.E. *see* Wotherspoon, F.
- Almdal, T. *see* Rosenfalck, A.M.
- Amiel, S.A. *see* Ashwell, S.G.
- Ammendola, C. *see* Marfella, R.
- Andersen, G. *see* Wegner, L.
- Anderson, J.M. *see* Porter, J.R.
- Anderson, R.A., Evans, L.M., Ellis, G.R., Khan, N., Morris, K., Jackson, S.K., Rees, A., Lewis, M.J. & Frenneaux, M.P. Prolonged deterioration of endothelial dysfunction in response to postprandial lipaemia is attenuated by vitamin C in Type 2 diabetes, 258
- Ando, A. *see* Muroi, R.
- Angelopoulos, N.G., Zervas, A., Livadas, S., Adamopoulos, I., Giannopoulos, D., Goula, A. & Tolis, G. Reduced insulin secretion in normoglycaemic patients with β -thalassaemia major, 1327
- Anguissola, G.B. *see* Pitocco, D.
- Anitha, B. *see* Rema, M.
- Anyaku, V. *see* Godsland, I.F.
- Araki, E. *see* Miyamura, N.
- Arcaro, G. *see* Targher, G.
- Argueta-Villamar, V. *see* Navalón-García, K.
- Arun, C.S., Young, D., Batey, D., Shotton, M., Mitchie, D., Stannard, K.P. & Taylor, R. Establishing ongoing quality assurance in a retinal screening programme, 629
- Ashwell, S.G., Amiel, S.A., Bilous, R.W., Dashora, U., Heller, S.R., Hepburn, D.A., Shutler, S.D., Stephens, J.W. & Home, P.D. Improved glycaemic control with insulin glargine plus insulin lispro: a multicentre, randomized, cross-over trial in people with Type 1 diabetes, 285
- Ashwell, S.G., Gebbie, J. & Home, P.D. Optimal timing of injection of once-daily insulin glargine in people with Type 1 diabetes using insulin lispro at meal-times, 46
- Ashwell, S.G., Gebbie, J. & Home, P.D. Twice-daily compared with once-daily insulin glargine in people with Type 1 diabetes using meal-time insulin aspart, 879
- Atay, A.E. *see* Ozmen, S.
- Atkins, M. *see* Linnebjerg, H.
- Attard, G. *see* Agius, E.
- Averous, C. *see* Kessler, L.
- Avierinos, J.-F. *see* Sejl, S.
- Avignon, A. *see* Sultan, A.
- Azizi, F. *see* Hosseiniapanah, F.
- Bacchetti, T. *see* Ferretti, G.
- Bader, L. *see* Petruzzo, P.
- Baek, H.S. *see* Park, J.H.
- Bahceci, M. *see* Ozmen, S.
- Bahr, C., Schön, S., Kuhn, J., Groop, P.H., Parkkonen, M., Wessman, M., Kleesiek, K. & Götting, C. Novel sequence variants in the human xylosyltransferase I gene and their role in diabetic nephropathy, 681

Author Index

- Aman, J. *see* Lodefalk, M.
- Abbas, J. *see* Forouhi, N.G.
- Abdella, N.A. *see* Mojiminiyi, O.A.
- Acerini, C.L. Weight matters for children. A complete guide for weight, eating and fitness, 1275
- Adaikalakoteswari, A., Balasubramanyam, M., Rema, M. & Mohan, V. Differential gene expression of NADPH oxidase (p22^{phox}) and hemoxygenase-1 in patients with Type 2 diabetes and microangiopathy, 666
- Adamopoulos, I. *see* Angelopoulos, N.G.
- Adang, E. *see* Dijkstra, R.F.
- Agarwal, M.M., Dhatt, G.S. & Punnose, J. Gestational diabetes: utility of fasting plasma glucose as a screening test depends on the diagnostic criteria, 1319
- Agius, E., Attard, G., Shakespeare, L., Clark, P., Vidya, M.-A., Hattersley, A.T. & Fava, S. Familial factors in diabetic nephropathy: an offspring study, 331
- Aguilar-Salinas, C.A. *see* Navalón-García, K.
- Ahmad, N. *see* Lawton, J.
- Ahmed, A.B. *see* Khan, A.S.A.
- Ahmed, A.T., Karter, A.J. & Liu, J. Alcohol consumption is inversely associated with adherence to diabetes self-care behaviours, 795
- Ahn, C.W. *see* Cho, N.H.
- Ahn, J.H., Kim, T.Y., Kim, Y.-J., Han, M.W., Yoon, T.H. & Chung, J.W. Lipo-prostaglandin E₁ in combination with steroid therapy is effective for treatment of sudden sensorineural hearing loss in Korean patients with Type 2 diabetes, 1339
- Ahn, T.Y. *see* Cho, N.H.
- Ajmani, A. *see* Vikram, N.K.
- Akram, K., Pedersen-Bjergaard, U., Carstensen, B., Borch-Johnsen, K. & Thorsteinsson, B. Frequency and risk factors of severe hypoglycaemia in insulin-treated Type 2 diabetes: a cross-sectional survey, 750
- Aksoy, M. *see* Ertugrul, M.B.
- Alberti, H. *see* Harzallah, F.
- Alberti, H. Diabetes in the Arab World, 107
- Alberti, K.G.M.M. *see* Whincup, P.
- Alberti, K.G.M.M., Zimmet, P. & Shaw, J. Metabolic syndrome—a new world-wide definition. A Consensus Statement from the International Diabetes Federation, 469
- Alberti, K.G.M.M., Zimmet, P. & Shaw, J. Reply, 1385
- Albrechtsen, A. *see* Wegner, L.
- Aldhahi, W.A. *see* Mojiminiyi, O.A.
- Alem, M. *see* Nazir, F.S.
- Ali, S., Stone, M.A., Peters, J.L., Davies, M.J. & Khunti, K. The prevalence of co-morbid depression in adults with Type 2 diabetes: a systematic review and meta-analysis, 1165
- Allard, S.E. *see* Browne, D.L.
- Allard, S.E. *see* Wotherspoon, F.
- Almdal, T. *see* Rosenfalck, A.M.
- Amiel, S.A. *see* Ashwell, S.G.
- Ammendola, C. *see* Marfella, R.
- Andersen, G. *see* Wegner, L.
- Anderson, J.M. *see* Porter, J.R.
- Anderson, R.A., Evans, L.M., Ellis, G.R., Khan, N., Morris, K., Jackson, S.K., Rees, A., Lewis, M.J. & Frenneaux, M.P. Prolonged deterioration of endothelial dysfunction in response to postprandial lipaemia is attenuated by vitamin C in Type 2 diabetes, 258
- Ando, A. *see* Muroi, R.
- Angelopoulos, N.G., Zervas, A., Livadas, S., Adamopoulos, I., Giannopoulos, D., Goula, A. & Tolis, G. Reduced insulin secretion in normoglycaemic patients with β -thalassaemia major, 1327
- Anguissola, G.B. *see* Pitocco, D.
- Anitha, B. *see* Rema, M.
- Anyaku, V. *see* Godsland, I.F.
- Araki, E. *see* Miyamura, N.
- Arcaro, G. *see* Targher, G.
- Argueta-Villamar, V. *see* Navalón-García, K.
- Arun, C.S., Young, D., Batey, D., Shotton, M., Mitchie, D., Stannard, K.P. & Taylor, R. Establishing ongoing quality assurance in a retinal screening programme, 629
- Ashwell, S.G., Amiel, S.A., Bilous, R.W., Dashora, U., Heller, S.R., Hepburn, D.A., Shutler, S.D., Stephens, J.W. & Home, P.D. Improved glycaemic control with insulin glargine plus insulin lispro: a multicentre, randomized, cross-over trial in people with Type 1 diabetes, 285
- Ashwell, S.G., Gebbie, J. & Home, P.D. Optimal timing of injection of once-daily insulin glargine in people with Type 1 diabetes using insulin lispro at meal-times, 46
- Ashwell, S.G., Gebbie, J. & Home, P.D. Twice-daily compared with once-daily insulin glargine in people with Type 1 diabetes using meal-time insulin aspart, 879
- Atay, A.E. *see* Ozmen, S.
- Atkins, M. *see* Linnebjerg, H.
- Attard, G. *see* Agius, E.
- Averous, C. *see* Kessler, L.
- Avierinos, J.-F. *see* Sejl, S.
- Avignon, A. *see* Sultan, A.
- Azizi, F. *see* Hosseiniapanah, F.
- Bacchetti, T. *see* Ferretti, G.
- Bader, L. *see* Petruzzo, P.
- Baek, H.S. *see* Park, J.H.
- Bahceci, M. *see* Ozmen, S.
- Bahr, C., Schön, S., Kuhn, J., Groop, P.H., Parkkonen, M., Wessman, M., Kleesiek, K. & Götting, C. Novel sequence variants in the human xylosyltransferase I gene and their role in diabetic nephropathy, 681

- Bailey, C. *see* Scanlon, P.H.
- Bailey, C.J. Fenofibrate and cardiovascular risk: a synopsis and commentary on (FIELD), 109
- Bain, S.C. *see* Dixon, A.N.
- Baker, J. *see* Kerr, G.D.
- Baktiroglu, S. *see* Ertugrul, M.B.
- Balasubramanyam, M. *see* Adaikalakoteswari, A.
- Balkau, B. *see* Cogneau, J.
- Bandello, F. *see* Trento, M.
- Banerjee, A.K. *see* Dixon, A.N.
- Barbieri, E. *see* Molon, G.
- Barnard, K.D., Skinner, T.C. & Peveler, R. The prevalence of comorbid depression in adults with Type 1 diabetes: systematic literature review, 445
- Baron, S. *see* Hazard, I.
- Barrett, T.G. *see* Porter, J.R.
- Baskar, V., Venugopal, H., Holland, M.R. & Singh, B.M. Clinical utility of estimated glomerular filtration rates in predicting renal risk in a district diabetes population, 1057
- Bassand, J.-P. *see* Schiele, F.
- Batey, D. *see* Arun, C.S.
- Batra, C.M. *see* Vikram, N.K.
- Bednarczuk, T. *see* Hiromatsu, Y.
- Begg, E.J. *see* Hughes, R.C.E.
- Ben Khalifa, F. *see* Harzallah, F.
- Benbow, S.J. *see* Daousi, C.
- Benecchi, R. *see* Trento, M.
- Berberoglu, K. *see* Ertugrul, M.B.
- Berger, B., Borg, H., Fernlund, P., Stenström, G. & Sundkvist, G. Islet antibodies associated with pancreatic B-cell dysfunction at and 3 years after diagnosis of diabetes in subjects aged 35–64 years old: degree of impairment less severe than in those aged 0–34 years old, 1180
- Berioli, M.G. *see* Celi, F.
- Berthillot, C. *see* Petruzzio, P.
- Berthold, H.K. *see* Gouni-Berthold, I.
- Bertolini, L. *see* Molon, G.
- Bertolini, L. *see* Targher, G.
- Bhattarai, M.D. *see* Shrestha, U.K.
- Bilous, R. Will ASCOT change the form of anti-hypertensive therapy?, 13
- Bilous, R.W. *see* Ashwell, S.G.
- Bilous, R.W. *see* New, J.P.
- Bingham, C. *see* Edghill, E.L.
- Bingley, P.J. *see* Castleden, H.A.J.
- Bini, V. *see* Celi, F.
- Biswas, N. *see* Stewart, M.W.
- Bizzarri, C. *see* Pitocco, D.
- Blaak, E.E. *see* Corpeleijn, E.
- Blakytyn, R. & Jude, E. The molecular biology of chronic wounds and delayed healing in diabetes, 594
- Blonde, M.C. *see* Schiele, F.
- Bobkiewicz-Kozłowska, T. *see* Dworacka, M.
- Bodansky, H.J. *see* Feltbower, R.G.
- Bodenlenz, M. *see* Schaller, H.C.
- Boeri, C. *see* Kessler, L.
- Bolli, G.B. *see* Hoogma, R.P.L.M.
- Booth, G.L. *see* Shah, B.R.
- Borch-Johnsen, K. *see* Akram, K.
- Borch-Johnsen, K. *see* Wegner, L.
- Borg, H. *see* Berger, B.
- Borodulin, K., Tuomilehto, J., Peltonen, M., Lakka, T.A., Sundvall, J. & Jousilahti, P. Association of leisure time physical activity and abdominal obesity with fasting serum insulin and 2-h postchallenge plasma glucose levels, 1025
- Boronat, M., Varillas, V.F., Saavedra, P., Suárez, V., Bosch, E., Carrillo, A. & Nóvoa, F.J. Diabetes mellitus and impaired glucose regulation in the Canary Islands (Spain): prevalence and associated factors in the adult population of Telde, Gran Canaria, 148
- Borowska, M. *see* Dworacka, M.
- Bosch, E. *see* Boronat, M.
- Bosio, A. *see* Fornengo, P.
- Bott, S., Tusek, C., Jacobsen, L.V., Endahl, L., Draeger, E., Kapitza, C. & Heise, T. Insulin detemir under steady-state conditions: no accumulation and constant metabolic effect over time with twice daily administration in subjects with Type 1 diabetes, 522
- Boucher, B.J. Hypovitaminosis D and risk of Type 2 diabetes in British South Asians, 336
- Boucher, B.J. Type 2 diabetes in rural and urban population: diverse prevalence and associated risk factors in Bangladesh, 450
- Bouhours-Nouet, N. *see* Hazard, I.
- Bouma, J. *see* Hagedoorn, M.
- Boumal, D. *see* Schiele, F.
- Bouter, K.P. *see* Hoogma, R.P.L.M.
- Bowden, D.W. *see* Burdon, K.P.
- Bowden, D.W., Lohman, K., Hsu, F.-C., Langefeld, C.D., Carr, J.J., Lenchik, L., Wagenknecht, L.E., Freedman, B.I. & Herrington, D.M. Hormone replacement therapy is associated with increased C-reactive protein in women with Type 2 diabetes in the Diabetes Heart Study, 763
- Bowker, S.L. *see* Johnson, J.A.
- Bradshaw, D. *see* Levitt, N.S.
- Brancato, R. *see* Trento, M.
- Braspenning, J.C.C. *see* Dijkstra, R.F.
- Brazert, J. *see* Dworacka, M.
- Brink, S. *see* Laffel, L.M.B.
- Brismar, K. *see* Ma, J.
- Brohall, G., Odén, A. & Fagerberg, B. Carotid artery intima-media thickness in patients with Type 2 diabetes mellitus and impaired glucose tolerance: a systematic review, 609
- Browne, D.L. *see* Wotherspoon, F.
- Browne, D.L., Meeking, D.R., Allard, S.E., Munday, J.L., Shaw, K.M. & Cummings, M.H. Angiotensin II does not affect endothelial tone in Type 1 diabetes—results of a double-blind placebo controlled trial, 53
- Brownstein, J.N. *see* Norris, S.L.
- Brunetti, P. *see* Perriello, G.
- Bruttomesso, D. *see* Hoogma, R.P.L.M.
- Bundy, C. *see* Knight, K.M.
- Burden, A.C. *see* Robertson, N.
- Burden, M.L. *see* Robertson, N.
- Burdge, G.C. *see* Masding, M.G.
- Burdon, K.P., Langefeld, C.D., Wagenknecht, L.E., Carr, J.J., Freedman, B.I., Herrington, D. & Bowden, D.W. Association analysis of genes in the renin-angiotensin system with subclinical cardiovascular disease in families with Type 2 diabetes mellitus: The Diabetes Heart Study, 228
- Busse Grawitz, A. *see* Slawik, M.

- Butler, R.A. *see* Franklin, V.L.
 Butzer, R. *see* Pfützner, J.
 Buxton, M.J. *see* Hanney, S.R.
 Buysschaert, M. *see* Hermans, M.P.
- Cacciapuoti, F. *see* Marfella, R.
 Cade, J.E. *see* Deakin, T.A.
 Calangu, S. *see* Ertugrul, M.B.
 Cameron, F.J. *see* Ismail, D.
 Cameron, F.J. *see* Northam, E.A.
 Campbell, F.M. *see* Feltbower, R.G.
 Canizales-Quinteros, S. *see* Navalón-García, K.
 Carr, J.J. *see* Bowden, D.W.
 Carr, J.J. *see* Burdon, K.P.
 Carrillo, A. *see* Boronat, M.
 Carroll, A.E. & Marrero, D.G. How do parents perceive their adolescent's diabetes: a qualitative study, 1222
 Carson, D.J. *see* Martin, R.J.L.
 Carstensen, B. *see* Akram, K.
 Casson, I.F. Pregnancy in women with diabetes—after the CEMACH report, what now?, 481
 Castellani, M.S. *see* Celi, F.
 Castleden, H.A.J., Shields, B., Bingley, P.J., Williams, A.J.K., Sampson, M., Walker, M., Gibson, J.M., McCarthy, M.I., Hitman, G.A., Levy, J.C., Hattersley, A.T., Vaidya, B. & Pearson, E.R. GAD antibodies in probands and their relatives in a cohort clinically selected for Type 2 diabetes, 834
 Cavallo, F. *see* Trento, M.
 Cavallo, M.G. *see* Pitocco, D.
 Cavan, D.A. *see* Shaban, M.C.
 Celi, F., Bini, V., Papi, F., Santilli, E., Castellani, M.S., Ferretti, A., Mencacci, M., Berlioli, M.G., De Giorgi, G. & Falorni, A. Circulating adipocytokines in non-diabetic and Type 1 diabetic children: relationship to insulin therapy, glycaemic control and pubertal development, 660
 Chan, J.C.N. *see* Yang, J.K.
 Chapman, P.G. *see* Johnson, N.A.
 Chassin, L.J. *see* Schaller, H.C.
 Chaturvedi, N. *see* Ferriss, J.B.
 Chen, X. *see* Liu, L.
 Cheong, H.S. *see* Park, K.S.
 Cheong, H.S. *see* Shin, H.D.
 Cheung, N.W. *see* Cutchie, W.A.
 Cheung, N.W. *see* McLean, M.
 Chien, J. *see* Wise, S.
 Chiorazzo, G. *see* Marfella, R.
 Chipps, D. *see* McLean, M.
 Cho, N.H., Ahn, C.W., Park, J.Y., Ahn, T.Y., Lee, H.W., Park, T.S., Kim, I.J., Pomerantz, K., Park, C., Kimm, K.C. & Choi, D.S. Prevalence of erectile dysfunction in Korean men with Type 2 diabetes mellitus, 198
 Cho, Y.M. *see* Park, K.S.
 Cho, Y.M. *see* Shin, H.D.
 Choi, D.S. *see* Cho, N.H.
 Chowdhury, F.M. *see* Norris, S.L.
 Chowdhury, T.A. *see* Moyes, V.
 Chowdhury, T.A. Reply, 1159
 Christmann, D. *see* Kessler, L.
 Chung, J.W. *see* Ahn, J.H.
 Cil, T. *see* Ozmen, S.
- Cinone, F. *see* Marfella, R.
 Cirkel, D.T. *see* Stewart, M.W.
 Civello, D. *see* Stock, S.A.K.
 Clark, P. *see* Agius, E.
 Cleland, S.J. *see* Nazir, F.S.
 Coggiola, A. *see* Trento, M.
 Cogneau, J., Balkau, B., Weill, A., Liard, F. & Simon, D. Assessment of diabetes screening by general practitioners in France: the EPIDIA Study, 803
 Colhoun, H.M. *see* Mulnier, H.E.
 Collober-Maugeais, C. *see* Ristic, S.
 Comte, F. *see* Sultan, A.
 Connell, J.M.C. *see* Nazir, F.S.
 Cook, D. *see* Whincup, P.
 Cook, N.R. *see* Pradhan, A.D.
 Coppini, D.V., Spruce, M.C., Thomas, P. & Masding, M.G. Established diabetic neuropathy seems irreversible despite improvements in metabolic and vascular risk markers—a retrospective case-control study in a hospital patient cohort, 1016
 Coppola, L. *see* Marfella, R.
 Cornel, M.C. *see* van Esch, S.C.M.
 Corpeleijn, E., van der Kallen, C.J.H., Kruijschoop, M., Magagnin, M.G.P., de Bruin, T.W.A., Feskens, E.J.M., Saris, W.H.M. & Blaak, E.E. Direct association of a promoter polymorphism in the CD36/FAT fatty acid transporter gene with Type 2 diabetes mellitus and insulin resistance, 907
 Corwin, P.A. *see* Gill, E.A.
 Costa, A. *see* Molon, G.
 Coutant, R. *see* Hazard, I.
 Craig, D. *see* Nelson, E.A.
 Craig, D. *see* O'Meara, S.
 Cressier, F. *see* Ristic, S.
 Crinò, A. *see* Pitocco, D.
 Croom, A. *see* Moyes, V.
 Crowle, T. *see* Sampson, M.J.
 Cummings, M.H. *see* Browne, D.L.
 Cummings, M.H. *see* Wotherspoon, F.
 Currie, C.J. *see* Morgan, C.L.I.
 Currie, C.J. Diabetes prevalence in England, 2001, 1268
 Cutchie, W.A., Cheung, N.W. & Simmons, D. Comparison of international and New Zealand guidelines for the care of pregnant women with diabetes, 460
- de la Calle, H. *see* Hoogma, R.P.L.M.
 de la Peña, A. *see* Linnebjerg, H.
 Diaz-Vargas, M.E. *see* Navalón-García, K.
 D'Andrea, F. *see* Marfella, R.
 Dahlquist, G. *see* Ma, J.
 Dai, M. *see* Jian, W.-X.
 Dalamanga, E.G. *see* Karamitsos, T.D.
 Dalton, J. *see* Nelson, E.A.
 Dalton, J.E. *see* O'Meara, S.
 Daly, M.E., Paisey, R., Paisey, R., Millward, B.A., Eccles, C., Williams, K., Hammersley, S., MacLeod, K.M. & Gale, T.J. Short-term effects of severe dietary carbohydrate-restriction advice in Type 2 diabetes—a randomized controlled trial, 15
 Damm, P. *see* Nielsen, L.R.
 Daousi, C., Benbow, S.J., Woodward, A. & MacFarlane, I.A. The natural history of chronic painful peripheral neuropathy in a community diabetes population, 1021

- Dasgupta, I. *see* New, J.P.
 Dashora, U. *see* Ashwell, S.G.
 Daures, J.P. *see* Sultan, A.
 Davidson, M., Zhu, J., Lu, W., Tracy, R.P., Robbins, D.C., Resnick, H.E., Ruotolo, G. & Howard, B.V., and the Strong Heart Study Investigators. Plasminogen activator inhibitor-1 and the risk of Type 2 diabetes mellitus in American Indians: the Strong Heart Study, 1158
 Davies, M., Dempster, M. & Malone, A. Do people with diabetes who need to talk want to talk?, 917
 Davies, M.J. *see* Ali, S.
 de Bruin, T.W.A. *see* Corpeleijn, E.
 de Fine Olivarius, N. & Jørgensen, L.G.M. Near-patient blood glucose measurements should only be used alone in the diagnosis of diabetes in cases of high glucose concentrations, 1042
 De Giorgi, G. *see* Celi, F.
 Deakin, T.A., Cade, J.E., Williams, R. & Greenwood, D.C. Structured patient education: the Diabetes X-PERT Programme makes a difference, 944
 Dean, J.D. *see* Young, B.
 Deary, I.J. *see* McAulay, V.
 Deepa, R. *see* Mohan, V.
 Deepa, R. *see* Rema, M.
 Delcourt, I. *see* Tauveron, I.
 Dempsey, E. *see* Gerstein, H.C.
 Dempster, M. *see* Davies, M.
 Desbiez, F. *see* Tauveron, I.
 Descotes-Genon, V. *see* Schiele, F.
 DeVane, J. *see* Jaffe, R.
 DeVries, J.H. Health-economic comparison of continuous subcutaneous insulin infusion with multiple daily injection for the treatment of Type 1 diabetes in the UK: Response to Roze et al., 709
 Dhatariya, K. *see* Sampson, M.J.
 Dhatt, G.S. *see* Agarwal, M.M.
 Di Pietro, C. *see* Perriello, G.
 Di Stasio, E. *see* Pitocco, D.
 Didangellos, T.P. *see* Karamitsos, T.D.
 Dijkstra, R.F., Niessen, L.W., Braspenning, J.C.C., Adang, E. & Grof, R.T.P.M. Patient-centred and professional-directed implementation strategies for diabetes guidelines: a cluster-randomized trial-based cost-effectiveness analysis, 164
 Dixon, A.N., Jude, E.B., Banerjee, A.K. & Bain, S.C. Simultaneous pulmonary and cerebral oedema, and multiple CNS infarctions as complications of diabetic ketoacidosis: a case report, 571
 Donaldson, J. *see* Stewart, M.W.
 Donath, S.M. *see* Ismail, D.
 Doney, A.S.F. *see* Donnelly, L.A.
 Donnelly, L.A., Doney, A.S.F., Hattersley, A.T., Morris, A.D. & Pearson, E.R. The effect of obesity on glycaemic response to metformin or sulphonylureas in Type 2 diabetes, 128
 Donovan, J. *see* Godslan, I.F.
 Dorel, S.B. *see* Petruzzo, P.
 Dornan, T. *see* Knight, K.M.
 Doughty, R.N. *see* Kerr, G.D.
 Douglas, M. *see* Lawton, J.
 Dozio, N. *see* Sampson, M.J.
 Draeger, E. *see* Bott, S.
 Driver, R. *see* Moyes, V.
 Duncan, E. *see* Toh, V.
 Duncan, G. *see* Shotliff, K.P.
 Dunseath, G. *see* Peter, R.
 Dworacka, M., Wender-Ozegowska, E., Winiarska, H., Borowska, M., Zawiejska, A., Pietryga, M., Brazert, J., Szczawinska, K. & Bobkiewicz-Kozłowska, T. Plasma anhydro-D-glucitol (1,5-AG) as an indicator of hyperglycaemic excursions in pregnant women with diabetes, 171
 Eagle, A.R.R. *see* Wolford, J.K.
 Ecartot, F. *see* Schiele, F.
 Eccles, C. *see* Daly, M.E.
 Edghill, E.L., Bingham, C., Slingerland, A.S., Minton, J.A.L., Noordam, C., Ellard, S. & Hattersley, A.T. Hepatocyte nuclear factor-1 beta mutations cause neonatal diabetes and intrauterine growth retardation: support for a critical role of HNF-1 β in human pancreatic development, 1301
 Edwards, A. *see* Johnson, J.A.
 Edwards, J. *see* Porter, J.R.
 Efendic, S. *see* Ma, J.
 Eher, R. *see* Maier, C.
 Eisenhut, M. Interleukin-1 and the constellation of pulmonary oedema, and cerebral infarctions and oedema in diabetic ketoacidosis, 1386
 El Gebely, S.A. *see* Mojiminiyi, O.A.
 Elkeles, R.S. *see* Godslan, I.F.
 Ellard, S. *see* Edghill, E.L.
 Ellard, S. *see* Porter, J.R.
 Ellepola, A.N.B. *see* Soysa, N.S.
 Ellis, G.R. *see* Anderson, R.A.
 Ellis, J.D. *see* Leese, G.P.
 Endahl, L. *see* Bott, S.
 English, D.R. *see* Hodge, A.M.
 Epifani, G. *see* Fornengo, P.
 Eriksson, J.G. *see* Mager, U.
 Ernest II, C.S. *see* Linnebjerg, H.
 Ertugrul, M.B., Baktiroglu, S., Salman, S., Unal, S., Aksoy, M., Berberoglu, K. & Calangu, S. The diagnosis of osteomyelitis of the foot in diabetes: microbiological examination vs. magnetic resonance imaging and labelled leucocyte scanning, 649
 Evans, L.M. *see* Anderson, R.A.
 Fagerberg, B. *see* Brohall, G.
 Falezza, G. *see* Targher, G.
 Falhammar, H. *see* Ma, J.
 Falorni, A. *see* Celi, F.
 Fang, Q. *see* Hu, C.
 Farjadian, S. *see* Mojtahedi, Z.
 Farmer, A., Kinmonth, A.-L. & Sutton, S. Measuring beliefs about taking hypoglycaemic medication among people with Type 2 diabetes, 265
 Faust, M. *see* Gouni-Berthold, I.
 Fava, S. *see* Agius, E.
 Feher, M.D. *see* Godslan, I.F.
 Felthower, R.G., Campbell, F.M., Bodansky, H.J., Stephenson, C.R. & McKinney, P.A. Insulin pump therapy in childhood diabetes—cost implications for Primary Care Trusts, 86
 Feng, Y. *see* Yang, J.K.
 Ferguson, B. *see* Forouhi, N.
 Ferguson, B.A. *see* Forouhi, N.G.

- Ferguson, S.C. & Frier, B.M. Hippocampal volume and cognitive performance in long-standing Type 1 diabetic patients without macrovascular complications, 1271
- Fernlund, P. *see* Berger, B.
- Ferretti, A. *see* Celi, F.
- Ferretti, G., Bacchetti, T., Rabini, R.A., Vignini, A., Nanetti, L., Moroni, C. & Mazzanti, L. Homocysteinylation of low-density lipoproteins (LDL) from subjects with Type 1 diabetes: effect on oxidative damage of human endothelial cells, 808
- Ferriss, J.B., Webb, D., Chaturvedi, N., Fuller, J.H. & Idzior-Walus, B.; The EURODIAB Prospective Complications Group. Weight gain is associated with improved glycaemic control but with adverse changes in plasma lipids and blood pressure in Type 1 diabetes, 557
- Feskens, E.J.M. *see* Corpeleijn, E.
- Fichter, L. *see* Toh, V.
- Fisher, B.M. *see* Mukhopadhyay, B.
- Fisher, M. Diabetes and Cardiovascular Disease, 929
- Flather, M.D. *see* Godsland, I.F.
- Fornengo, P., Bosio, A., Epifani, G., Pallisco, O., Mancuso, A. & Pascale, C. Prevalence of silent myocardial ischaemia in new-onset middle-aged Type 2 diabetic patients without other cardiovascular risk factors, 775
- Forouhi, N., Goyder, E., Merrick, D., Wild, S. & Ferguson, B. Authors' reply, 1268
- Forouhi, N.G. *see* Mukhopadhyay, B.
- Forouhi, N.G., Merrick, D., Goyder, E., Ferguson, B.A., Abbas, J., Lachowycz, K. & Wild, S.H. Diabetes prevalence in England, 2001—estimates from an epidemiological model, 189
- Forsblom, C. *see* Turunen, J.A.
- Forst, S. *see* Pfützner, J.
- Forst, T. *see* Pfützner, J.
- Fosbury, J. *see* Shaban, M.C.
- Frame, I. *see* Hanney, S.R.
- Franklin, V.L., Waller, A., Pagliari, C. & Greene, S.A. A randomized controlled trial of Sweet Talk, a text-messaging system to support young people with diabetes, 1332
- Franklin, V.L., Wilson, A.W., Butler, R.A. & Greene, S.A. A predictive tool for the self-management of diabetes (Librae): evaluation using a continuous glucose monitoring system, 21
- Fraser, L. *see* Ismail, D.
- Frayling, T.M. *see* Porter, J.R.
- Freed, M.I. *see* Viberti, G.
- Freedman, B.I. *see* Bowden, D.W.
- Freedman, B.I. *see* Burdon, K.P.
- Frenneaux, M.P. *see* Anderson, R.A.
- Frier, B.M. *see* Ferguson, S.C.
- Frier, B.M. *see* Geddes, J.
- Frier, B.M. *see* McAulay, V.
- Fu, H.J. *see* Yang, J.K.
- Fujita, N. *see* Muroi, R.
- Fukutani, T. *see* Hiromatsu, Y.
- Fuller, J.H. *see* Ferriss, J.B.
- Furuse, K. *see* Stewart, M.W.
- Götting, C. *see* Bahr, C.
- Gaboury, I. *see* Malcolm, J.C.
- Gala, J.-L. *see* Hermans, M.P.
- Gale, T.J. *see* Daly, M.E.
- Gall, M.-A. *see* Kølendorf, K.
- Gamble, G.D. *see* Kerr, G.D.
- Gardiner, S.J. *see* Hughes, R.C.E.
- Garvey, W.T. *see* Lyons, T.J.
- Gatelais, F. *see* Hazard, I.
- Gaudias, J. *see* Kessler, L.
- Gebbie, J. *see* Ashwell, S.G.
- Gebert, R. *see* Ismail, D.
- Geddes, J., McGeough, E. & Frier, B.M. Young adults with Type 1 diabetes in tertiary education: do students receive adequate specialist care?, 1155
- Gerhard-Herman, M.D. *see* Raji, A.
- Gerstein, H.C., Yale, J.-F., Harris, S.B., Issa, M., Stewart, J.A. & Dempsey, E. A randomized trial of adding insulin glargine vs. avoidance of insulin in people with Type 2 diabetes on either no oral glucose-lowering agents or submaximal doses of metformin and/or sulphonylureas. The Canadian INSIGHT (Implementing New Strategies with Insulin Glargine for Hyperglycaemia Treatment) Study, 736
- Ghaderi, A. *see* Mojtahedi, Z.
- Giannakidou, E. *see* Gouni-Berthold, I.
- Giannopoulos, D. *see* Angelopoulos, N.G.
- Gibson, J.M. *see* Castleden, H.A.J.
- Giles, G.G. *see* Hodge, A.M.
- Gilg, J. *see* Whincup, P.
- Gill, E.A., Corwin, P.A., Mangin, D.A. & Sutherland, M.G. Diabetes care in rest homes in Christchurch, New Zealand, 1252
- Gill, G. *see* Woodward, A.
- Giorda, C., Petrelli, A. & Gnani, R., the Regional Board for Diabetes Care of Piemonte. The impact of second-level specialized care on hospitalization in persons with diabetes: a multilevel population-based study, 377
- Glümer, C. *see* Wegner, L.
- Gloyn, A.L. *see* Porter, J.R.
- Gnani, R. *see* Giorda, C.
- Godsland, I.F., Elkeles, R.S., Feher, M.D., Nugara, F., Rubens, M.B., Richmond, W., Khan, M., Donovan, J., Anyaoku, V. & Flather, M.D., for the PREDICT Study Group. Coronary calcification, homocysteine, C-reactive protein and the metabolic syndrome in Type 2 diabetes: the Prospective Evaluation of Diabetic Ischaemic Heart Disease by Coronary Tomography (PREDICT) Study, 1192
- Goh, S.K. *see* Lim, S.C.
- Gokulakrishnan, K. *see* Mohan, V.
- Golder, S. *see* Nelson, E.A.
- Golder, S. *see* O'Meara, S.
- Gomis, R. *see* Hoogma, R.P.L.M.
- González-Chávez, A. *see* Navalón-García, K.
- Gough, S.C. *see* Palin, S.L.
- Goula, A. *see* Angelopoulos, N.G.
- Gouni-Berthold, I., Giannakidou, E., Faust, M., Berthold, H.K. & Krone, W. Association of the promoter polymorphism -232C/G of the phosphoenolpyruvate carboxykinase gene (PCK1) with Type 2 diabetes mellitus, 419
- Goyder, E. *see* Forouhi, N.
- Goyder, E. *see* Forouhi, N.G.
- Gram, J. *see* Heldgaard, P.E.
- Grant, I. *see* Nyirenda, M.J.
- Grant, J. *see* Hanney, S.R.

- Gravenor, M.B. *see* Pegge, N.C.
- Green, P. *see* Hanney, S.R.
- Greene, S.A. *see* Franklin, V.L.
- Greenwood, D.C. *see* Deakin, T.A.
- Greenwood, R.H. *see* Sampson, M.J.
- Griffin, S.J. *see* Heldgaard, P.E.
- Grol, R.T.P.M. *see* Dijkstra, R.F.
- Groop, P.H. *see* Bahr, C.
- Groop, P.H. *see* Turunen, J.A.
- Gu, H.F. *see* Ma, J.
- Gu, Y.-Y. *see* Jian, W.-X.
- Guglielmi, C. *see* Pitocco, D.
- Gupta, N. *see* Vikram, N.K.
- Häger, S. *see* von Sengbusch, S.
- Hämäläinen, H. *see* Mager, U.
- Haan, M.N. *see* Pradhan, A.D.
- Habib, G. *see* Sejil, S.
- Haffner, S. *see* Viberti, G.
- Hagedoorn, M., Keers, J.C., Links, T.P., Bouma, J., Ter Maaten, J.C. & Sanderma, R. Improving self-management in insulin-treated adults participating in diabetes education. The role of overprotection by the partner, 271
- Hallowell, N. *see* Lawton, J.
- Hamada, Y. *see* Watarai, A.
- Hamaguchi, T. *see* Satani, K.
- Hamajima, N. *see* Watarai, A.
- Hamann, A. *see* Stewart, M.W.
- Hammersley, S. *see* Daly, M.E.
- Hammond, P.J. *see* Hoogma, R.P.L.M.
- Han, B.G. *see* Park, K.S.
- Han, B.G. *see* Shin, H.D.
- Han, J.-F. *see* Jian, W.-X.
- Han, M.W. *see* Ahn, J.H.
- Hanifi-Moghaddam, P., Kappler, S., Seissler, J., Müller-Scholz, S., Martin, S., Roep, B.O., Strassburger, K., Kolb, H. & Schloot, N.C. Altered chemokine levels in individuals at risk of Type 1 diabetes mellitus, 156
- Hanna, L. *see* Lawton, J.
- Hanney, S.R., Home, P.D., Frame, I., Grant, J., Green, P. & Buxton, M.J. Identifying the impact of diabetes research, 176
- Hansen, T. *see* Wegner, L.
- Hansmann, Y. *see* Kessler, L.
- Hanson, R.L. *see* Wolford, J.K.
- Hara, T. *see* Tsutsumi, Y.
- Harper, C.A. *see* Phiri, R.
- Harris, S.B. *see* Gerstein, H.C.
- Harten-Ash, V. Repeat education with self-monitoring of blood glucose improves HbA_{1c}, 220
- Harzallah, F., Alberti, H. & Ben Khalifa, F. The metabolic syndrome in an Arab population: a first look at the new International Diabetes Federation criteria, 441
- Hasegawa, Y. *see* Satani, K.
- Hatazaki, M. *see* Nakatani, Y.
- Hattersley, A.T. *see* Agius, E.
- Hattersley, A.T. *see* Castleden, H.A.J.
- Hattersley, A.T. *see* Donnelly, L.A.
- Hattersley, A.T. *see* Edghill, E.L.
- Hattersley, A.T. *see* Porter, J.R.
- Hattersley, A.T. *see* Stenhouse, E.
- Hawthorne, G. Metformin use and diabetic pregnancy—has its time come?, 223
- Hazard, I., Bouhours-Nouet, N., Baron, S., Ramos, E., Gatelais, F. & Coutant, R. Birthweight predicts insulin requirements in children and adolescents with Type 1 diabetes, 1273
- He, K. *see* Song, Y.
- Hearnshaw, H. & Lindenmeyer, A. What do we mean by adherence to treatment and advice for living with diabetes? A review of the literature on definitions and measurements, 720
- Heise, M.A. *see* Viberti, G.
- Heise, T. *see* Bott, S.
- Heldgaard, P.E. & Griffin, S.J. Routinely collected general practice data aids identification of people with hyperglycaemia and metabolic syndrome, 996
- Heldgaard, P.E., Sidelmann, J.J., Hindsberger, C., Olivarius, N. de F., Henriksen, J.E. & Gram, J. Relationship of glucose concentrations with PAI-1 and t-PA in subjects with normal glucose tolerance, 887
- Heller, S.R. *see* Ashwell, S.G.
- Heller, S.R. *see* Kolendorf, K.
- Hendrix, S.L. *see* Pradhan, A.D.
- Henriksen, J.E. *see* Heldgaard, P.E.
- Hepburn, D.A. *see* Ashwell, S.G.
- Herman, W.H. *see* Viberti, G.
- Hermans, M.P., Gala, J.-L. & Buysschaert, M. The MTHFR C₆₇₇T polymorphism confers a high risk for stroke in both homozygous and heterozygous T allele carriers with Type 2 diabetes, 529
- Herrington, D. *see* Burdon, K.P.
- Herrington, D.M. *see* Bowden, D.W.
- Heyburn, P.J. *see* Sampson, M.J.
- Hilsted, J. *see* Rosenfalck, A.M.
- Hindsberger, C. *see* Heldgaard, P.E.
- Hiort, O. *see* von Sengbusch, S.
- Hiromatsu, Y., Mukai, T., Kaku, H., Miyake, I., Ichimura, M., Fukutani, T., Nakayama, H., Takata, K., Imamura, Y., Shoji, S., Yamada, K., Koda, Y. & Bednarczyk, T. IL-18 gene polymorphism confers susceptibility to the development of anti-GAD65 antibody in Graves' disease, 211
- Hitchman, R. *see* McElduff, A.
- Hitman, G.A. *see* Castleden, H.A.J.
- Hjemdahl, P. *see* Yngen, M.
- Hodge, A.M., English, D.R., O'Dea, K. & Giles, G.G. Alcohol intake, consumption pattern and beverage type, and the risk of Type 2 diabetes, 690
- Holland, M.R. *see* Baskar, V.
- Holland, P. *see* Milton, B.
- Holman, R. *see* Viberti, G.
- Home, P.D. *see* Ashwell, S.G.
- Home, P.D. *see* Hanney, S.R.
- Home, P.D. Cost-effectiveness of CSII, 707
- Hoogma, R.P.L.M., Hammond, P.J., Gomis, R., Kerr, D., Bruttomesso, D., Bouter, K.P., Wiefels, K.J., de la Calle, H., Schweitzer, D.H., Pfohl, M., Torlone, E., Krinkel, L.G. & Bolli, G.B., on behalf of the 5-Nations Study Group. Comparison of the effects of continuous subcutaneous insulin infusion (CSII) and NPH-based multiple daily insulin injections (MDI) on glycaemic control and quality of life: results of the 5-nations trial, 141
- Horsley, T. *see* Norris, S.L.

- Hosseiniapanah, F., Rambod, M., Reza Ghaffari, H.R. & Azizi, F. Predicting isolated postchallenge hyperglycaemia: a new approach; Tehran Lipid and Glucose Study (TLGS), 982
- Hovorka, R. *see* Schaller, H.C.
- Hovorka, R. Continuous glucose monitoring and closed-loop systems, 1
- Howard, B.V. *see* Davidson, M.
- Hsu, F.-C. *see* Bowden, D.W.
- Hu, C., Jia, W., Fang, Q., Zhang, R., Wang, C., Lu, J. & Xiang, K. *Peroxisome proliferator-activated receptor (PPAR) delta* genetic polymorphism and its association with insulin resistance index and fasting plasma glucose concentrations in Chinese subjects, 1307
- Huang, C. *see* Liu, L.
- Huang, G. *see* Zhang, D.
- Hughes, R.C.E. & Rowan, J.A. Pregnancy in women with Type 2 diabetes: who takes metformin and what is the outcome?, 318
- Hughes, R.C.E., Gardiner, S.J., Begg, E.J. & Zhang, M. Effect of pregnancy on the pharmacokinetics of metformin, 323
- Hundborg, H.H. *see* Kruse, A.
- Hussain, A. Reply, 450
- Hux, J.E. *see* Shah, B.R.
- Hykin, P. *see* Scanlon, P.H.
- Ichihara, Y. *see* Miyamura, N.
- Ichimura, M. *see* Hiromatsu, Y.
- Ideno, J. *see* Muroi, R.
- IDF Clinical Guidelines Task Force. Global Guideline for Type 2 Diabetes: recommendations for standard, comprehensive, and minimal care, 579
- Idzior-Walus, B. *see* Ferriss, J.B.
- Iglesias, C. *see* Nelson, E.A.
- Iglesias, C. *see* O'Meara, S.
- Ihara, K. *see* Tsutsumi, Y.
- Ilanne-Parikka, P. *see* Mager, U.
- Imamura, Y. *see* Hiromatsu, Y.
- Insacco, C. *see* Trento, M.
- Ishay, A., Lavi, I. & Luboshitzky, R. Prevalence and risk factors for asymptomatic bacteriuria in women with Type 2 diabetes mellitus, 185
- Ishibashi, S. *see* Muroi, R.
- Ishikawa, T. *see* Satani, K.
- Ismail, D., Gebert, R., Vuillermin, P.J., Fraser, L., McDonnell, C.M., Donath, S.M. & Cameron, F.J. Social consumption of alcohol in adolescents with Type 1 diabetes is associated with increased glucose lability, but not hypoglycaemia, 830
- Ismail, H., Wright, J., Rhodes, P. & Scally, A. Quality of care in diabetic patients attending routine primary care clinics compared with those attending GP specialist clinics, 851
- Issa, M. *see* Gerstein, H.C.
- Jørgensen, L.G.M. *see* de Fine Olivarius, N.
- Jørgensen, T. *see* Wegner, L.
- Jack Jr, L. *see* Norris, S.L.
- Jackson, S.K. *see* Anderson, R.A.
- Jacobsen, L.V. *see* Bott, S.
- Jacobsen, P. *see* Lajer, M.
- Jaffe, R., Mani, J., DeVane, J. & Mani, H. Tolerance loss in diabetics: association with foreign antigen exposure, 924
- Jakobsen, P. *see* Mortensen, U.M.
- James, W.H. The sex ratios of offspring of diabetic parents, 1043
- Janand-Delenne, B. *see* Seijl, S.
- Janghorbani, M., Stenhouse, E.A., Jones, R.B. & Millward, B.A. Is neighbourhood deprivation a risk factor for gestational diabetes mellitus?, 313
- Jeffcoate, W.J. The evidence base to guide the use of antibiotics in foot ulcers in people with diabetes is thin, but what are we going to do about it?, 339
- Jendle, J. *see* Kølendorf, K.
- Jenkins, A.J. *see* Lyons, T.J.
- Jensen, B.S. *see* Mortensen, U.M.
- Jerums, G. *see* Thomas, M.C.
- Jia, W. *see* Hu, C.
- Jian, W.-X., Luo, T.-H., Gu, Y.-Y., Zhang, H.-L., Zheng, S., Dai, M., Han, J.-F., Zhao, Y., Li, G. & Luo, M. The visfatin gene is associated with glucose and lipid metabolism in a Chinese population, 967
- Jie, X. *see* Tsutsumi, Y.
- Jin, F. *see* Sun, L.
- Jing, P. *see* Zhang, D.
- Johnson, E. *see* Scanlon, P.H.
- Johnson, J.A., Majumdar, S.R., Bowker, S.L., Toth, E.L. & Edwards, A. Self-monitoring in Type 2 diabetes: a randomized trial of reimbursement policy, 1247
- Johnson, K.C. *see* Pradhan, A.D.
- Johnson, N.A., Stannard, S.R., Rowlands, D.S., Chapman, P.G., Thompson, C.H., Sachinwalla, T. & Thompson, M.W. Short-term suppression of plasma free fatty acids fails to improve insulin sensitivity when intramyocellular lipid is elevated, 1061
- Jones, C. *see* Sampson, M.J.
- Jones, N.P. *see* Viberti, G.
- Jones, R.B. *see* Janghorbani, M.
- Jousilahti, P. *see* Borodulin, K.
- Jude, E. *see* Blakytyn, R.
- Jude, E.B. *see* Dixon, A.N.
- Jung, R.T. *see* Leese, G.P.
- Kølendorf, K., Ross, G.P., Pavlic-Renar, I., Perriello, G., Philotheou, A., Jendle, J., Gall, M.-A. & Heller, S.R. Insulin detemir lowers the risk of hypoglycaemia and provides more consistent plasma glucose levels compared with NPH insulin in Type 1 diabetes, 729
- Kahn, S.E. *see* Viberti, G.
- Kakishita, E. *see* Satani, K.
- Kaku, H. *see* Hiromatsu, Y.
- Kamiya, H. *see* Watarai, A.
- Kanemitsu, S. *see* Tsutsumi, Y.
- Kaneto, H. *see* Nakatani, Y.
- Kapitza, C. *see* Bott, S.
- Kappler, S. *see* Hanifi-Moghaddam, P.
- Karamitsos, D.T. *see* Karamitsos, T.D.
- Karamitsos, T.D., Karvounis, H.I., Didangellos, T.P., Papadopoulos, C.E., Dalamanga, E.G., Karamitsos, D.T., Parharidis, G.E. & Louridas, G.E. Usefulness of colour tissue Doppler imaging in assessing aortic elastic properties in Type 1 diabetic patients, 1201
- Karter, A.J. *see* Ahmed, A.T.
- Karvounis, H.I. *see* Karamitsos, T.D.
- Kasper, J. *see* Lenz, M.
- Katsuno, T. *see* Satani, K.

- Kawamori, D. *see* Nakatani, Y.
 Keeffe, J.E. *see* Phiri, R.
 Keely, E. *see* Malcolm, J.C.
 Keers, J.C. *see* Hagedoorn, M.
 Keightley, S. *see* Scanlon, P.H.
 Keinänen-Kiukaanniemi, S. *see* Mager, U.
 Kerr, D. & Richardson, T. Treating insulin allergy with continuous subcutaneous insulin infusion (CSII), 1159
 Kerr, D. *see* Hoogma, R.P.L.M.
 Kerr, D. *see* Shaban, M.C.
 Kerr, G.D., Gamble, G.D., Doughty, R.N., Simmons, D. & Baker, J. Mortality in individuals with Type 2 diabetes and heart disease in a unique New Zealand population, 1313
 Kessler, L., Piemont, Y., Ortega, F., Lesens, O., Boeri, C., Averous, C., Meyer, R., Hansmann, Y., Christmann, D., Gaudias, J. & Pinget, M. Comparison of microbiological results of needle puncture vs. superficial swab in infected diabetic foot ulcer with osteomyelitis, 99
 Kesson, C.M. *see* Mukhopadhyay, B.
 Khan, A.S.A., McLoughney, C.R. & Ahmed, A.B. The effect of metformin on blood glucose control in overweight patients with Type 1 diabetes, 1079
 Khan, M. *see* Godslan, I.F.
 Khan, N. *see* Anderson, R.A.
 Khunti, K. *see* Ali, S.
 Kilpikari, R. *see* Turunen, J.A.
 Kim, H.T. *see* Park, K.S.
 Kim, H.T. *see* Shin, H.D.
 Kim, I.J. *see* Cho, N.H.
 Kim, J.H. *see* Park, J.H.
 Kim, J.W. *see* Shin, H.D.
 Kim, T.Y. *see* Ahn, J.H.
 Kim, W.H. *see* Park, J.H.
 Kim, Y.-J. *see* Ahn, J.H.
 Kim, Y.J. *see* Shin, H.D.
 Kimm, K. *see* Park, K.S.
 Kimm, K. *see* Shin, H.D.
 Kimm, K.C. *see* Cho, N.H.
 Kinmonth, A.-L. *see* Farmer, A.
 Kleesiek, K. *see* Bahr, C.
 Klein, R.L. *see* Lyons, T.J.
 Knight, K.M., Dornan, T. & Bundy, C. The diabetes educator: trying hard, but must concentrate more on behaviour, 485
 Knowler, W.C. *see* Wolford, J.K.
 Knudsen, L.L. *see* Kruse, A.
 Kobayashi, H. *see* Muroi, R.
 Kobayashi, Y. *see* Watarai, A.
 Koda, Y. *see* Hiromatsu, Y.
 Koh, A. *see* Lim, S.C.
 Koh, I. *see* Shin, H.D.
 Kohri, K. *see* Satani, K.
 Kolb, H. *see* Hanifi-Moghaddam, P.
 Kolm, P. *see* Rhee, M.K.
 Konya, H. *see* Satani, K.
 Kothare, P.A. *see* Linnebjerg, H.
 Krömeke, O. *see* Lobnig, B.M.
 Kravitz, B. *see* Viberti, G.
 Krinkel, L.G. *see* Hoogma, R.P.L.M.
 Krone, W. *see* Gouni-Berthold, I.
 Kruijschoop, M. *see* Corpeleijn, E.
 Kruse, A., Thomsen, R.W., Hundborg, H.H., Knudsen, L.L., Sørensen, H.T. & Schönheyder, H.C. Diabetes and risk of acute infectious conjunctivitis—a population-based case-control study, 393
 Kuhn, J. *see* Bahr, C.
 Kuo, T.B.J. *see* Sun, T.B.
 Laakso, M. *see* Mager, U.
 Labastie, N. *see* Sejl, S.
 Lachin, J. *see* Viberti, G.
 Lachowycz, K. *see* Forouhi, N.G.
 Lackland, D.T. *see* Lyons, T.J.
 LaCroix, A.Z. *see* Pradhan, A.D.
 Laffel, L.M.B., Wentzell, K., Loughlin, C., Tovar, A., Moltz, K. & Brink, S. Sick day management using blood 3-hydroxybutyrate (3-OHB) compared with urine ketone monitoring reduces hospital visits in young people with T1DM: a randomized clinical trial, 278
 Lai, M.-Y. *see* Su, D.-H.
 Lai, Y.R. *see* Lim, S.C.
 Laight, D.W. *see* Wotherspoon, F.
 Lajer, M., Schjoedt, K.J., Jacobsen, P., Tarnow, L. & Parving, H.-H. Aldosterone synthase (CYP11B2) -344T/C polymorphism is not associated with the initiation and progression of diabetic nephropathy in Caucasian Type 1 diabetic patients, 675
 Lakka, T.A. *see* Borodulin, K.
 Langefeld, C.D. *see* Bowden, D.W.
 Langefeld, C.D. *see* Burdon, K.P.
 Lassmann-Vague, V. *see* Sejl, S.
 Lattanzio, R. *see* Trento, M.
 Laupacis, A. *see* Shah, B.R.
 Lauterbach, K.W. *see* Stock, S.A.K.
 Lavi, I. *see* Ishay, A.
 Laville, M. *see* Petruzzio, P.
 Lawrenson, R.A. *see* Mulnier, H.E.
 Lawson, M.L. *see* Malcolm, J.C.
 Lawton, J., Ahmad, N., Hanna, L., Douglas, M. & Hallowell, N. Diabetes service provision: a qualitative study of the experiences and views of Pakistani and Indian patients with Type 2 diabetes, 1003
 Leary, S.D. *see* Shultis, W.A.
 Lee, B.L. *see* Lim, S.C.
 Lee, H.K. *see* Park, K.S.
 Lee, H.K. *see* Shin, H.D.
 Lee, H.W. *see* Cho, N.H.
 Lee, J.-K. *see* Park, K.S.
 Lee, J.-K. *see* Shin, H.D.
 Lee, J.-Y. *see* Park, K.S.
 Lee, J.-Y. *see* Shin, H.D.
 Lees, K.R. *see* Nazir, F.S.
 Leese, G.P., Morris, A.D., Petrie, J., Jung, R.T. & Ellis, J.D. Reply, 449
 Lefrançois, N. *see* Petruzzio, P.
 Legalery, P. *see* Schiele, F.
 Lenchik, L. *see* Bowden, D.W.
 Lenz, M., Kasper, J. & Mühlhauser, I. Searching for diabetes decision aids and related background information, 912
 Lesens, O. *see* Kessler, L.
 Leslie, E. *see* Xu, F.
 Levitan, E.B. *see* Song, Y.

- Levitt, N.S. & Bradshaw, D. The impact of HIV/AIDS on Type 2 diabetes prevalence and diabetes healthcare needs in South Africa: projections for 2010, 103
- Levy, J.C. *see* Castleden, H.A.J.
- Lewis, M.J. *see* Anderson, R.A.
- Lewis, N. *see* Toh, V.
- Li, G. *see* Jian, W.-X.
- Li, L. *see* Zhang, D.
- Li, X. *see* Sundquist, K.
- Liard, F. *see* Cogneau, J.
- Liberti, D. *see* Marfella, R.
- Lim, S.C., Goh, S.K., Lai, Y.R., Tee, W.W., Koh, A., Xu, X.H., Wu, Y.S., Yap, E., Subramaniam, T. & Sum, C.F. Relationship between common functional polymorphisms of the p22phox gene (-930A > G and +242C > T) and nephropathy as a result of Type 2 diabetes in a Chinese population, 1037
- Lim, S.C., Tan, H.H., Goh, S.K., Subramaniam, T., Sum, C.F., Tan, I.K., Lee, B.L. & Ong, C.N. Oxidative burden in prediabetic and diabetic individuals: evidence from plasma coenzyme Q₁₀, 1344
- Lindenmeyer, A. *see* Hearnshaw, H.
- Lindi, V. *see* Mager, U.
- Lindström, J. *see* Mager, U.
- Links, T.P. *see* Hagedoorn, M.
- Linnebjerg, H., Kothare, P.A., Skrivaneck, Z., de la Peña, A., Atkins, M., Ernest II, C.S. & Trautmann, M.E. Exenatide: effect of injection time on postprandial glucose in patients with Type 2 diabetes, 240
- Liu, J. *see* Ahmed, A.T.
- Liu, L., Yu, Q., Wang, H., Zhang, S.X., Huang, C. & Chen, X. Association of intercellular adhesion molecule 1 polymorphisms with retinopathy in Chinese patients with Type 2 diabetes, 643
- Liu, S. *see* Song, Y.
- Livadas, S. *see* Angelopoulos, N.G.
- Lobnig, B.M. & Wolf, O.T. Reply, 1272
- Lobnig, B.M., Krömeke, O., Optenhostert-Porst, C. & Wolf, O.T. Hippocampal volume and cognitive performance in long-standing Type 1 diabetic patients without macrovascular complications, 32
- Lodefalk, M. & Åman, J. Food habits, energy and nutrient intake in adolescents with Type 1 diabetes mellitus, 1225
- Lohman, K. *see* Bowden, D.W.
- Lopes-Virella, M. *see* Lyons, T.J.
- Lough, G. *see* Malcolm, J.C.
- Loughlin, C. *see* Laffel, L.M.B.
- Louridas, G.E. *see* Karamitsos, T.D.
- Lu, J. *see* Hu, C.
- Lu, W. *see* Davidson, M.
- Luboshitzky, R. *see* Ishay, A.
- Lucassen, P.L.B.J. *see* van de Laar, F.A.
- Luger, A. *see* Maier, C.
- Luo, M. *see* Jian, W.-X.
- Luo, T.-H. *see* Jian, W.-X.
- Luzio, S.D. *see* Peter, R.
- Lyons, T.J., Jenkins, A.J., Zheng, D., Klein, R.L., Otvos, J.D., Yu, Y., Lackland, D.T., McGee, D., McHenry, M.B., Lopes-Virella, M. & Garvey, W.T., The DCCT/EDIC Research Group. Nuclear magnetic resonance-determined lipoprotein subclass profile in the DCCT/EDIC cohort: associations with carotid intima-media thickness, 955
- Möllsten, A. *see* Ma, J.
- Mühlhauser, I. *see* Lenz, M.
- Müller, C. *see* Nielsen, L.R.
- Müller-Godeffroy, E. *see* von Sengbusch, S.
- Müller-Scholze, S. *see* Hanifi-Moghaddam, P.
- Ma, J., Möllsten, A., Prázný, M., Falhammar, H., Brismar, K., Dahlquist, G., Efendic, S. & Gu, H.F. Genetic influences of the intercellular adhesion molecule 1 (ICAM-1) gene polymorphisms in development of Type 1 diabetes and diabetic nephropathy, 1093
- MacFarlane, I.A. *see* Daousi, C.
- MacIsaac, R. *see* Thomas, M.C.
- MacLeod, K.M. *see* Daly, M.E.
- Madhu, S.V. *see* Vikram, N.K.
- Madsbad, S. *see* Rosenfalk, A.M.
- Magagnin, M.G.P. *see* Corpeleijn, E.
- Mager, U., Lindi, V., Lindström, J., Eriksson, J.G., Valle, T.T., Hämäläinen, H., Ilanne-Parikka, P., Keinänen-Kiukaanniemi, S., Tuomilehto, J., Laakso, M., Pulkkinen, L. & Uusitupa, M. for the Finnish Diabetes Prevention Study Group. Association of the Leu72Met polymorphism of the ghrelin gene with the risk of Type 2 diabetes in subjects with impaired glucose tolerance in the Finnish Diabetes Prevention Study, 685
- Maier, C., Mustapic, D., Schuster, E., Luger, A. & Eher, R. Effect of a pocket-size tablet-dispensing device on glycaemic control in Type 2 diabetic patients, 40
- Majumdar, S.R. *see* Johnson, J.A.
- Malcolm, J.C., Lawson, M.L., Gaboury, I., Lough, G. & Keely, E. Glucose tolerance of offspring of mother with gestational diabetes mellitus in a low-risk population, 565
- Malone, A. *see* Davies, M.
- Malone, J.K. *see* Roach, P.
- Mancuso, A. *see* Fornengo, P.
- Manfrini, S. *see* Pitocco, D.
- Mangin, D.A. *see* Gill, E.A.
- Mani, H. *see* Jaffe, R.
- Mani, J. *see* Jaffe, R.
- Manson, J.E. *see* Pradhan, A.D.
- Manson, J.E. *see* Song, Y.
- Marchese, F. *see* Marfella, R.
- Marfella, M.A. *see* Marfella, R.
- Marfella, R., Cacciapuoti, F., Siniscalchi, M., Sasso, F.C., Marchese, F., Cinone, F., Musacchio, E., Marfella, M.A., Ruggiero, L., Chiorazzo, G., Liberti, D., Chiorazzo, G., Nicoletti, G.F., Saron, C., D'Andrea, F., Ammendola, C., Verza, M. & Coppola, L. Effect of moderate red wine intake on cardiac prognosis after recent acute myocardial infarction of subjects with Type 2 diabetes mellitus, 974
- Mariano-Goulart, D. *see* Sultan, A.
- Marrero, D.G. *see* Carroll, A.E.
- Marshall, S.M. *see* New, J.P.
- Martínez-Godínez, M.A. *see* Navalón-García, K.
- Martin, M.L. *see* Scanlon, P.H.
- Martin, R.J.L., Savage, D.A., Carson, D.J., Maxwell, A.P. & Patterson, C.C. The PAX4 gene variant A1168C is not associated with early onset Type 1 diabetes in a UK population, 927
- Martin, R.M. *see* Shultis, W.A.
- Martin, S. *see* Hanifi-Moghaddam, P.
- Martin, X. *see* Petruzzio, P.
- Masding, M.G. *see* Coppini, D.V.

- Masding, M.G., Stears, A.J., Burdge, G.C., Wootton, S.A. & Sandeman, D.D. The benefits of oestrogens on postprandial lipid metabolism are lost in post-menopausal women with Type 2 diabetes, 768
- Mathiesen, E.R. *see* Nielsen, L.R.
- Matsuhisa, M. *see* Nakatani, Y.
- Matsumoto, K. *see* Miyamura, N.
- Matsumura, T. *see* Miyamura, N.
- Matsuoka, T. *see* Nakatani, Y.
- Matteoli, M.C. *see* Pitocco, D.
- Matthews, D.R. *see* Toh, V.
- Matyka, K. Clinical Obesity in Adults and Children, 1160
- Maxwell, A.P. *see* Martin, R.J.L.
- Mazzanti, L. *see* Ferretti, G.
- McAulay, V., Deary, I.J., Sommerfield, A.J. & Frier, B.M. Attentional functioning is impaired during acute hypoglycaemia in people with Type 1 diabetes, 26
- McCarthy, M.I. *see* Castleden, H.A.J.
- McDonnell, C.M. *see* Ismail, D.
- McElduff, A., Hitchman, R. & McElduff, P. Is sex hormone-binding globulin associated with glucose tolerance?, 306
- McElduff, P. *see* McElduff, A.
- McEwan, P. *see* Morgan, C.L.L.
- McGee, D. *see* Lyons, T.J.
- McGeough, E. *see* Geddes, J.
- McHenry, M.B. *see* Lyons, T.J.
- McKinney, P.A. *see* Feltbower, R.G.
- McKnight, J.A. *see* Nyirenda, M.J.
- McLean, M., Chipps, D. & Cheung, N.W. Mother to child transmission of diabetes mellitus: does gestational diabetes program Type 2 diabetes in the next generation?, 1213
- McLoughney, C.R. *see* Khan, A.S.A.
- Meeking, D.R. *see* Browne, D.L.
- Meeking, D.R. *see* Wotherspoon, F.
- Mencacci, M. *see* Celi, F.
- Mendoza-Alcantar, L. *see* Navalón-García, K.
- Meneveau, N. *see* Schiele, F.
- Mercier, M. *see* Schiele, F.
- Merrick, D. *see* Forouhi, N.
- Merrick, D. *see* Forouhi, N.G.
- Meyer, R. *see* Kessler, L.
- Middleton, R.J. *see* New, J.P.
- Miliar-García, A. *see* Navalón-García, K.
- Millward, B.A. *see* Daly, M.E.
- Millward, B.A. *see* Janghorbani, M.
- Millward, B.A. *see* Stenhouse, E.
- Milton, B., Holland, P. & Whitehead, M. The social and economic consequences of childhood-onset Type 1 diabetes mellitus across the lifecourse: a systematic review, 821
- Minton, J.A.L. *see* Edghill, E.L.
- Mirakian, R. *see* Moyes, V.
- Misra, A. *see* Vikram, N.K.
- Mitchie, D. *see* Arun, C.S.
- Mithal, A. *see* Vikram, N.K.
- Miwa, K. *see* Watarai, A.
- Miyajima, H. *see* Muroi, R.
- Miyake, I. *see* Hiromatsu, Y.
- Miyamura, N., Murata, Y., Taketa, K., Ichihara, Y., Matsumura, T., Tokunaga, H., Matsumoto, K., Sakakida, M. & Araki, E. A case of insulin autoimmune syndrome with HLA DRB1*0404: impact on the hypothesis for the molecular pathogenesis involving DRB1 molecules, 104
- Mohamed, H.M. *see* Mojiminiyi, O.A.
- Mohan, V. *see* Adaikalakoteswari, A.
- Mohan, V. *see* Rema, M.
- Mohan, V., Gokulakrishnan, K., Sandeep, S., Srivastava, B.K., Ravikumar, R. & Deepa, R. Intimal media thickness, glucose intolerance and metabolic syndrome in Asian Indians—the Chennai Urban Rural Epidemiology Study (CURES-22), 845
- Mojiminiyi, O.A., Abdella, N.A., Zaki, M.Y., El Gebely, S.A., Mohamed, H.M. & Aldhahi, W.A. Prevalence and associations of low plasma erythropoietin in patients with Type 2 diabetes mellitus, 839
- Mojtahedi, Z., Naeimi, S., Farjadian, S., Omrani, G.R. & Ghaderi, A. Association of IL-18 promoter polymorphisms with predisposition to Type 1 diabetes, 235
- Molon, G., Targher, G., Costa, A., Bertolini, L., Barbieri, E. & Zenari, L. Measurement of microvolt T-wave alternans, a new arrhythmic risk stratification test, in Type 2 diabetic patients without clinical cardiovascular disease, 207
- Moltz, K. *see* Laffel, L.M.B.
- Montanaro, M. *see* Trento, M.
- Morgan, C.L.L., McEwan, P., Morrissey, M., Peters, J.R., Poole, C. & Currie, C.J. Characterization and comparison of health-related utility in people with diabetes with various single and multiple vascular complications, 1100
- Mori, M. *see* Shimizu, H.
- Moroni, C. *see* Ferretti, G.
- Morris, A.D. *see* Donnelly, L.A.
- Morris, A.D. *see* Leese, G.P.
- Morris, K. *see* Anderson, R.A.
- Morrissey, M. *see* Morgan, C.L.L.
- Mortensen, U.M., Nielsen-Kudsk, J.E., Sondergaard, H.M., Jakobsen, P., Jensen, B.S., Schmitz, O. & Nielsen, T.T. Effect of the oral hypoglycaemic sulphonylurea glibenclamide, a blocker of ATP-sensitive potassium channels, on walking distance in patients with intermittent claudication, 327
- Moyes, V., Driver, R., Croom, A., Mirakian, R. & Chowdhury, T.A. Insulin allergy in a patient with Type 2 diabetes successfully treated with continuous subcutaneous insulin infusion, 204
- Mukai, T. *see* Hiromatsu, Y.
- Mukhopadhyay, B., Forouhi, N.G., Fisher, B.M., Kesson, C.M. & Sattar, N. A comparison of glycaemic and metabolic control over time among South Asian and European patients with Type 2 diabetes: results from follow-up in a routine diabetes clinic, 94
- Mulder, J. *see* van de Laar, F.A.
- Mulnier, H.E., Seaman, H.E., Raleigh, V.S., Soedamah-Muthu, S.S., Colhoun, H.M. & Lawrenson, R.A. Mortality in people with Type 2 diabetes in the UK, 516
- Munday, J.L. *see* Browne, D.L.
- Munday, L.J. *see* Wotherspoon, F.
- Murata, Y. *see* Miyamura, N.
- Muroi, R., Yagyu, H., Kobayashi, H., Nagata, M., Sato, N., Ideno, J., Fujita, N., Ando, A., Okada, K., Takiyama, Y., Nagasaka, S., Miyajima, H., Nakano, I. & Ishibashi, S. Early onset insulin-dependent diabetes mellitus as an initial manifestation of aceruloplasminemia, 1136
- Murphy, H.R., Rayman, G. & Skinner, T.C. Psycho-educational interventions for children and young people with Type 1 diabetes, 935

- Musacchio, E. *see* Marfella, R.
 Mustafa, N. *see* Peter, R.
 Mustapic, D. *see* Maier, C.
- Nóvoa, F.J. *see* Boronat, M.
 Nacimi, S. *see* Mojtaheidi, Z.
 Nagasaka, S. *see* Muroi, R.
 Nagata, M. *see* Muroi, R.
 Nakae, M. *see* Watarai, A.
 Nakamura, J. *see* Watarai, A.
 Nakano, I. *see* Muroi, R.
 Nakashima, E. *see* Watarai, A.
 Nakatani, Y., Kaneto, H., Hatazaki, K., Yoshiuchi, K., Kawamori, D., Sakamoto, K., Matsuoka, T., Ogawa, S., Yamasaki, Y. & Matsuhisa, M. Increased stress protein ORP150 autoantibody production in Type 1 diabetic patients, 216
 Nakayama, H. *see* Hiromatsu, Y.
 Namba, M. *see* Satani, K.
 Nanetti, L. *see* Ferretti, G.
 Naruse, K. *see* Watarai, A.
 Navalón-García, K., Mendoza-Alcantar, L., Díaz-Vargas, M.E., Martínez-Godínez, M.A., Reyna-Garfias, H., Aguilar-Salinas, C.A., Riba, L., Canizales-Quinteros, S., Villarreal-Molina, T., González-Chávez, A., Argueta-Villamar, V., Tusié-Luna, M.T. & Miliar-García, A. HNF-1 α G574S is a functional variant with decreased transactivation activity, 1295
 Nazir, F.S., Alem, M., Small, M., Connell, J.M.C., Lees, K.R., Walters, M.R. & Cleland, S.J. Blunted response to systemic nitric oxide synthase inhibition in the cerebral circulation of patients with Type 2 diabetes, 398
 Nelson, E.A. *see* O'Meara, S.
 Nelson, E.A., O'Meara, S., Golder, S., Dalton, J., Craig, D. & Iglesias, C., on behalf of the DASIDU Steering Group. Systematic review of antimicrobial treatments for diabetic foot ulcers, 348
 Nelson, R.G. *see* Wolford, J.K.
 Ness, A.R. *see* Shults, W.A.
 New, J.P., O'Donoghue, D.J., Middleton, R.J., Rudenski, A., Dasgupta, I., Bilous, R.W. & Marshall, S.M. Time to move from serum creatinine to eGFR, 1047
 Nicoletti, G.F. *see* Marfella, R.
 Nielsen, L.R., Müller, C., Damm, P. & Mathiesen, E.R. Reduced prevalence of early preterm delivery in women with Type 1 diabetes and microalbuminuria—possible effect of early antihypertensive treatment during pregnancy, 426
 Nielsen, T.T. *see* Mortensen, U.M.
 Nielsen-Kudsk, J.E. *see* Mortensen, U.M.
 Niessen, L.W. *see* Dijkstra, R.F.
 Niwa, T. *see* Watarai, A.
 Nomura, A. *see* Tsutsumi, Y.
 Noordam, C. *see* Edghill, E.L.
 Norris, S.L., Chowdhury, F.M., Van Le, K., Horsley, T., Brownstein, J.N., Zhang, X., Jack Jr, L. & Satterfield, D.W. Effectiveness of community health workers in the care of persons with diabetes, 544
 Northam, E.A., Todd, S. & Cameron, F.J. Interventions to promote optimal health outcomes in children with Type 1 diabetes—are they effective?, 113
 Nugara, F. *see* Godsland, I.F.
- Nyirenda, M.J., Sandeep, T., Grant, I., Price, G. & McKnight, J.A. Severe acidosis in patients taking metformin—rapid reversal and survival despite high APACHE score, 432
- Östenson, C.-G. *see* Yngen, M.
 O'Dea, K. *see* Hodge, A.M.
 O'Donoghue, D.J. *see* New, J.P.
 O'Meara, S. *see* Nelson, E.A.
 O'Meara, S., Nelson, E.A., Golder, S., Dalton, J.E., Craig, D. & Iglesias, C., on behalf of the DASIDU Steering Group. Systematic review of methods to diagnose infection in foot ulcers in diabetes, 341
 O'Connor, M.E. *see* Raji, A.
 O'Neill, M.C. *see* Viberti, G.
 Odén, A. *see* Brohall, G.
 Oda, E. Cut points of waist circumference for Japanese people, 1385
 Ogawa, S. *see* Nakatani, Y.
 Oh, B. *see* Park, K.S.
 Oh, B. *see* Shin, H.D.
 Oh-I, S. *see* Shimizu, H.
 Ohtani, K.-I. *see* Shimizu, H.
 Oiso, Y. *see* Watarai, A.
 Okada, K. *see* Muroi, R.
 Okada, S. *see* Shimizu, H.
 Olivarius, N. de F. *see* Heldgaard, P.E.
 Omrani, G.R. *see* Mojtaheidi, Z.
 Ong, C.N. *see* Lim, S.C.
 Optenhostert-Porst, C. *see* Lobnig, B.M.
 Ortega, F. *see* Kessler, L.
 Orvos, J.D. *see* Lyons, T.J.
 Owen, C. *see* Whincup, P.
 Owen, N. *see* Xu, F.
 Owens, D.R. *see* Peter, R.
 Ozmen, S., Cil, T., Atay, A.E., Tuzcu, A.K. & Bahceci, M. A simple way to estimate mean plasma glucose and to identify Type 2 diabetic subjects with poor glycaemic control when a standardized HbA_{1c} assay is not available, 1151
- Padovani, R. *see* Targher, G.
 Pagliari, C. *see* Franklin, V.L.
 Paisey, R. *see* Daly, M.E.
 Palin, S.L. & Gough, S.C. Rupture of the Achilles tendon associated with ciprofloxacin, 1386
 Pallisco, O. *see* Fornengo, P.
 Palmer, A.J. *see* Roze, S.
 Pampanelli, S. *see* Perriello, G.
 Pandey, R.M. *see* Vikram, N.K.
 Papadopoulos, C.E. *see* Karamitsos, T.D.
 Papi, F. *see* Celi, F.
 Parharidis, G.E. *see* Karamitsos, T.D.
 Park, B.L. *see* Park, K.S.
 Park, B.L. *see* Shin, H.D.
 Park, C. *see* Cho, N.H.
 Park, C. *see* Shin, H.D.
 Park, C.-S. *see* Park, K.S.
 Park, J.H., Kim, W.H., Kim, J.H., Park, T.S. & Baek, H.S. Prevalence of and risk factors for extracranial internal carotid artery stenosis in Korean Type 2 diabetic patients, 1377
 Park, J.Y. *see* Cho, N.H.
 Park, K.S. *see* Shin, H.D.

- Park, K.S., Shin, H.D., Park, B.L., Cheong, H.S., Cho, Y.M., Lee, H.K., Lee, J.-Y., Lee, J.-K., Kim, H.T., Park, C.-S., Han, B.G., Kimm, K. & Oh, B. Putative association of peroxisome proliferator-activated receptor γ co-activator 1 β (*PPARGC1B*) polymorphism with Type 2 diabetes mellitus, 635
- Park, T.S. *see* Cho, N.H.
- Park, T.S. *see* Park, J.H.
- Parkkonen, M. *see* Bahr, C.
- Parkkonen, M. *see* Turunen, J.A.
- Parving, H.-H. *see* Lajer, M.
- Pascale, C. *see* Fornengo, P.
- Patera, I.P. *see* Pitocco, D.
- Patterson, C.C. *see* Martin, R.J.L.
- Pauvaday, V. *see* Peter, R.
- Pavlic-Renar, I. *see* Kølendorf, K.
- Pearson, E.R. *see* Castleden, H.A.J.
- Pearson, E.R. *see* Donnelly, L.A.
- Pecher, E. *see* Ristic, S.
- Pedersen, O. *see* Wegner, L.
- Pedersen-Bjergaard, U. *see* Akram, K.
- Pegge, N.C., Twomey, A.M., Vaughton, K., Gravenor, M.B., Ramsey, M.W. & Price, D.E. The role of endothelial dysfunction in the pathophysiology of erectile dysfunction in diabetes and in determining response to treatment, 873
- Peltonen, M. *see* Borodulin, T.
- Penforis, A. *see* Schiele, F.
- Peng, J. *see* Zhang, D.
- Perriello, G. *see* Kølendorf, K.
- Perriello, G., Pampanelli, S., Di Pietro, C. & Brunetti, P., on behalf of the Italian Pioglitazone Study Group. Comparison of glycaemic control over 1 year with pioglitazone or gliclazide in patients with Type 2 diabetes, 246
- Peter, R., Luzio, S.D., Dunseath, G., Pauvaday, V., Mustafa, N. & Owens, D.R. Relationship between HbA_{1c} and indices of glucose tolerance derived from a standardized meal test in newly diagnosed treatment naive subjects with Type 2 diabetes, 990
- Peters, J.L. *see* Ali, S.
- Peters, J.R. *see* Morgan, C.L.I.
- Petersen, K.-G. *see* Slawik, M.
- Petrelli, A. *see* Giorda, C.
- Petrie, J. *see* Leese, G.P.
- Petruzzio, P., Badet, L., Lefrançois, N., Berthillot, C., Dorel, S.B., Martin, X. & Laville, M. Metabolic consequences of pancreatic systemic or portal venous drainage in simultaneous pancreas-kidney transplant recipients, 654
- Peveler, R. *see* Barnard, K.D.
- Pfützner, A. *see* Pfützner, J.
- Pfützner, A.H. *see* Pfützner, J.
- Pfützner, J., Forst, T., Butzer, R., Forst, S., Weber, M.M., Pfützner, A.H. & Pfützner, A. Performance of the continuous glucose monitoring system (CGMS) during development of ketosis in patients on insulin pump therapy, 1124
- Pfohl, M. *see* Hoogma, R.P.L.M.
- Phenikos, C. *see* Stewart, M.W.
- Phillips, L.S. *see* Rhee, M.K.
- Philothou, A. *see* Kølendorf, K.
- Phiri, R., Keefe, J.E., Harper, C.A. & Taylor, H.R. Comparative study of the polaroid and digital non-mydiatic cameras in the detection of refractive diabetic retinopathy in Australia, 867
- Pieber, T.R. *see* Schaller, H.C.
- Piemont, Y. *see* Kessler, L.
- Pietryga, M. *see* Dworacka, M.
- Pinget, M. *see* Kessler, L.
- Piot, C. *see* Sultan, A.
- Pitocco, D., Crinò, A., Di Stasio, E., Manfrini, S., Guglielmi, C., Spera, S., Anguissola, G.B., Visalli, N., Suraci, C., Matteoli, M.C., Patera, I.P., Cavallo, M.G., Bizzarri, C. & Pozzilli, P., on behalf of the IMDIAB Group. The effects of calcitriol and nicotinamide on residual pancreatic β -cell function in patients with recent-onset Type 1 diabetes (IMDIAB XI), 920
- Plummer, C.J. What's in the CARDS?, 711
- Plunkett, M. *see* Porter, J.R.
- Poli, F. *see* Targher, G.
- Pomerantz, K. *see* Cho, N.H.
- Poole, C. *see* Morgan, C.L.I.
- Porta, M. *see* Trento, M.
- Porter, J.R., Rangasami, J.J., Ellard, S., Gloyn, A.L., Shields, B.M., Edwards, J., Anderson, J.M., Shaw, N.J., Hattersley, A.T., Frayling, T.M., Plunkett, M. & Barrett, T.G. Asian MODY: are we missing an important diagnosis?, 1257
- Pozzilli, P. *see* Pitocco, D.
- Prázný, M. *see* Ma, J.
- Pradhan, A.D., Manson, J.E., Hendrix, S.L., Johnson, K.C., Wagenknecht, L.E., Haan, M.N., Weidner, G., LaCroix, A.Z. & Cook, N.R. Cross-sectional correlates of fasting hyperinsulinaemia in post-menopausal women of different ethnic origin, 77
- Price, D.E. *see* Pegge, N.C.
- Price, G. *see* Nyirenda, M.J.
- Pulkkinen, L. *see* Mager, U.
- Punnose, J. *see* Agarwal, M.M.
- Qu, Y.C. *see* Sun, L.
- Rabini, R.A. *see* Ferretti, G.
- Racah, D. *see* Sevil, S.
- Raji, A., Gerhard-Herman, M.D., Williams, J.S., O'Connor, M.E. & Simonson, D.C. Effect of pioglitazone on insulin sensitivity, vascular function and cardiovascular inflammatory markers in insulin-resistant non-diabetic Asian Indians, 537
- Raleigh, V.S. *see* Mulnier, H.E.
- Rambod, M. *see* Hosseinpahan, F.
- Ramos, E. *see* Hazard, I.
- Ramsey, M.W. *see* Pegge, N.C.
- Rangasami, J.J. *see* Porter, J.R.
- Ravikumar, R. *see* Mohan, V.
- Rayman, G. *see* Murphy, H.R.
- Redaelli, M. *see* Stock, S.A.K.
- Rees, A. *see* Anderson, R.A.
- Reincke, M. *see* Slawik, M.
- Reintjes, R. *see* von Sengbusch, S.
- Rema, M. *see* Adaikalakoteswari, A.
- Rema, M., Srivastava, B.K., Anitha, B., Deepa, R. & Mohan, V. Association of serum lipids with diabetic retinopathy in urban South Indians—the Chennai Urban Rural Epidemiology Study (CURES) Eye Study—2, 1029
- Renard, E. *see* Sultan, A.
- Rendas-Baum, R. *see* Roy, M.
- Resnick, H.E. *see* Davidson, M.
- Reyna-Garfias, H. *see* Navalón-García, K.

- Reza Ghaffari, H.R. *see* Hosseiniapanah, F.
- Rhee, M.K., Ziemer, D.C., Kolm, P. & Phillips, L.S. Postchallenge glucose rises with increasing age even when glucose tolerance is normal, 1174
- Rhodes, P. *see* Ismail, H.
- Riba, L. *see* Navalón-García, K.
- Richardson, C. *see* Wise, S.
- Richardson, T. *see* Kerr, D.
- Richmond, W. *see* Godsland, I.F.
- Ristic, S., Collober-Maugeais, C., Pecher, E. & Cressier, F.
Comparison of nateglinide and gliclazide in combination with metformin, for treatment of patients with Type 2 diabetes mellitus inadequately controlled on maximum doses of metformin alone, 757
- Roach, P. & Malone, J.K. Comparison of insulin lispro mixture 25/75 with insulin glargine during a 24-h standardized test-meal period in patients with Type 2 diabetes, 743
- Robbins, D.C. *see* Davidson, M.
- Robertson, N., Burden, M.L. & Burden, A.C. Psychological morbidity and problems of daily living in people with visual loss and diabetes: do they differ from people without diabetes?, 1110
- Roep, B.O. *see* Hanifi-Moghaddam, P.
- Rooney, D.P. & Ryan, M.F. Diabetes with partial lipodystrophy following sclerodermatous chronic graft vs. host disease, 436
- Rosenfalk, A.M., Almdal, T., Viggers, L., Madsbad, S. & Hilsted, J. A low-fat diet improves peripheral insulin sensitivity in patients with Type 1 diabetes, 384
- Ross, G.P. *see* Kølendorf, K.
- Rowan, J.A. *see* Hughes, R.C.E.
- Rowlands, D.S. *see* Johnson, N.A.
- Roy, M., Rendas-Baum, R. & Skurnick, J. Mortality in African-Americans with Type 1 diabetes: The New Jersey 725, 698
- Roze, S., Valentine, W.J., Zakrzewska, K. & Palmer, A.J. Response to the letter by Philip Home, 707
- Rubens, M.B. *see* Godsland, I.F.
- Rudenski, A. *see* New, J.P.
- Ruggiero, L. *see* Marfella, R.
- Ruotolo, G. *see* Davidson, M.
- Rutten, G.E.H.M. *see* van de Laar, F.A.
- Ryan, M.F. *see* Rooney, D.P.
- Sørensen, H.T. *see* Kruse, A.
- Saavedra, P. *see* Boronat, M.
- Sachinwalla, T. *see* Johnson, N.A.
- Sakakida, M. *see* Miyamura, N.
- Sakamoto, K. *see* Nakatani, Y.
- Salman, S. *see* Ertugrul, M.B.
- Samaranayake, L.P. *see* Soysa, N.S.
- Sampson, M. *see* Castleden, H.A.J.
- Sampson, M.J., Crowle, T., Dhatariya, K., Dozio, N., Greenwood, R.H., Heyburn, P.J., Jones, C., Temple, R.C. & Walden, E. Trends in bed occupancy for inpatients with diabetes before and after the introduction of a diabetes inpatient specialist nurse service, 1008
- Sandbaek, A. Reply, 1042
- Sandeep, S. *see* Mohan, V.
- Sandeep, T. *see* Nyirenda, M.J.
- Sandeman, D.D. *see* Masding, M.G.
- Sanderman, R. *see* Hagedoorn, M.
- Santilli, E. *see* Celi, F.
- Saris, W.H.M. *see* Corpeleijn, E.
- Saron, C. *see* Marfella, R.
- Sasso, F.C. *see* Marfella, R.
- Satani, K., Konya, H., Hamaguchi, T., Umehara, A., Katsuno, T., Ishikawa, T., Kohri, K., Hasegawa, Y., Suehiro, A., Kakishita, E. & Namba, M. Clinical significance of circulating hepatocyte growth factor, a new risk marker of carotid atherosclerosis in patients with Type 2 diabetes, 617
- Sato, N. *see* Muroi, R.
- Sattar, N. *see* Mukhopadhyay, B.
- Satterfield, D.W. *see* Norris, S.L.
- Savage, D.A. *see* Martin, R.J.L.
- Scala, L. *see* Targher, G.
- Scally, A. *see* Ismail, H.
- Scanlon, P.H., Martin, M.L., Bailey, C., Johnson, E., Hykin, P. & Keightley, S. Reported symptoms and quality-of-life impacts in patients having laser treatment for sight-threatening diabetic retinopathy, 60
- Schönheyder, H.C. *see* Kruse, A.
- Schön, S. *see* Bahr, C.
- Schaller, H.C., Schaupp, L., Bodenlenz, M., Wilinska, M.E., Chassin, L.J., Wach, P., Vering, T., Hovorka, R. & Pieber, T.R. On-line adaptive algorithm with glucose prediction capacity for subcutaneous closed loop control of glucose: evaluation under fasting conditions in patients with Type 1 diabetes, 90
- Schaupp, L. *see* Schaller, H.C.
- Schiele, F., Descotes-Genon, V., Seronde, M.F., Blonde, M.C., Legall, P., Meneveau, N., Ecarnot, F., Mercier, M., Penfornis, A., Thebault, L., Boumal, D. & Bassand, J-P., on behalf of the investigators of the 'Réseau Franc Comtois de Cardiologie'. Predictive value of admission hyperglycaemia on mortality in patients with acute myocardial infarction, 1370
- Schjoedt, K.J. *see* Lajer, M.
- Schlott, N.C. *see* Hanifi-Moghaddam, P.
- Schmitz, O. *see* Mortensen, U.M.
- Schories, M. *see* Slawik, M.
- Schuster, E. *see* Maier, C.
- Schweitzer, D.H. *see* Hoogma, R.P.L.M.
- Scoccianti, L. *see* Trento, M.
- Scott, J. *see* Shultis, W.A.
- Scottish Study Group for the Care of the Young with Diabetes. A longitudinal observational study of insulin therapy and glycaemic control in Scottish children with Type 1 diabetes: DIAB³UD 3, 1216
- Seaman, H.E. *see* Mulnier, H.E.
- Seissler, J. *see* Hanifi-Moghaddam, P.
- Sejil, S., Janand-Delenne, B., Avierinos, J.-F., Habib, G., Labastie, N., Raccach, D., Vague, P. & Lassmann-Vague, V. Six-year follow-up of a cohort of 203 patients with diabetes after screening for silent myocardial ischaemia, 1186
- Sekido, Y. *see* Watarai, A.
- Seronde, M.F. *see* Schiele, F.
- Shaban, M.C., Fosbury, J., Kerr, D. & Cavan, D.A. The prevalence of depression and anxiety in adults with Type 1 diabetes, 1381
- Shah, B.R., Hux, J.E., Laupacis, A., Zinman, B. & Booth, G.L. Use of vascular risk-modifying medications for diabetic patients differs between physician specialties, 1117
- Shakespeare, L. *see* Agius, E.
- Sharma, S. *see* Vikram, N.K.

- Shashaj, B. *see* Sulli, N.
- Shaw, J. *see* Alberti, K.G.M.M.
- Shaw, K.M. *see* Browne, D.L.
- Shaw, K.M. *see* Wotherspoon, F.
- Shaw, N.J. *see* Porter, J.R.
- Shi, X.H. *see* Sun, L.
- Shields, B. *see* Castleden, H.A.J.
- Shields, B.M. *see* Porter, J.R.
- Shimizu, H., Oh-I, S., Tsuchiya, T., Ohtani, K.-I., Okada, S. & Mori, M. Pioglitazone increases circulating adiponectin levels and subsequently reduces TNF- α levels in Type 2 diabetic patients: a randomized study, 253
- Shin, H.D. *see* Park, K.S.
- Shin, H.D., Park, K.S., Park, B.L., Cheong, H.S., Cho, Y.M., Lee, H.K., Lee, J.-Y., Lee, J.-K., Kim, H.T., Han, B.G., Kim, J.W., Koh, I., Kim, Y.J., Oh, B., Kimm, K. & Park, C. Common promoter polymorphism in monocyte differentiation antigen CD14 is associated with serum triglyceride levels and body mass index in non-diabetic individuals, 72
- Shoji, S. *see* Hiromatsu, Y.
- Shotliff, K.P. & Duncan, G. Diabetic retinopathy screening programmes and reducing ophthalmologists' workload, 449
- Shotton, M. *see* Arun, C.S.
- Shrestha, U.K., Singh, D.L. & Bhattarai, M.D. The prevalence of hypertension and diabetes defined by fasting and 2-h plasma glucose criteria in urban Nepal, 1130
- Shultis, W.A., Leary, S.D., Ness, A.R., Scott, J., Martin, R.M., Whincup, P.H. & Smith, G.D., the ALSPAC Study Team. Haemoglobin A_{1c} is not a surrogate for glucose and insulin measures for investigating the early life and childhood determinants of insulin resistance and Type 2 diabetes in healthy children. An analysis from the Avon Longitudinal Study of Parents and Children (ALSPAC), 1357
- Shutler, S.D. *see* Ashwell, S.G.
- Sidelmann, J.J. *see* Heldgaard, P.E.
- Simmons, D. *see* Cutchie, W.A.
- Simmons, D. *see* Kerr, G.D.
- Simon, D. *see* Cogneau, J.
- Simonson, D.C. *see* Raji, A.
- Singh, B.M. *see* Baskar, V.
- Singh, D.L. *see* Shrestha, U.K.
- Siniscalchi, M. *see* Marfella, R.
- Skinner, T.C. *see* Barnard, K.D.
- Skinner, T.C. *see* Murphy, H.R.
- Skinner, T.C. What does make the difference?, 933
- Skrivanek, Z. *see* Linnebjerg, H.
- Skurnick, J. *see* Roy, M.
- Slawik, M., Schories, M., Busse Grawitz, A., Reincke, M. & Petersen, K.-G. Treatment with insulin glargine does not suppress serum IGF-1, 814
- Slingerland, A.S. *see* Edghill, E.L.
- Small, M. *see* Nazir, F.S.
- Smith, G.D. *see* Shultis, W.A.
- Snoek, F.J. *see* van Esch, S.C.M.
- Soedamah-Muthu, S.S. *see* Mulnier, H.E.
- Somda, F. *see* Tauveron, I.
- Sommerfield, A.J. *see* McAulay, V.
- Sondergaard, H.M. *see* Mortensen, U.M.
- Song, Y., He, K., Levitan, E.B., Manson, J.E. & Liu, S. Effects of oral magnesium supplementation on glycaemic control in Type 2 diabetes: a meta-analysis of randomized double-blind controlled trials, 1050
- Soysa, N.S., Samaranyake, L.P. & Ellepola, A.N.B. Diabetes mellitus as a contributory factor in oral candidosis, 455
- Sparso, T. *see* Wegner, L.
- Spera, S. *see* Pitocco, D.
- Spruce, M.C. *see* Coppini, D.V.
- Srivastava, B.K. *see* Mohan, V.
- Srivastava, B.K. *see* Rema, M.
- Srivastava, M.C. *see* Vikram, N.K.
- Stafleu, A. *see* van de Laar, F.A.
- Stannard, K.P. *see* Arun, C.S.
- Stannard, S.R. *see* Johnson, N.A.
- Starkie, M.G. *see* Stewart, M.W.
- Stears, A.J. *see* Masding, M.G.
- Stenhouse, E., Wright, D.E., Hattersley, A.T. & Millward, B.A. Maternal glucose levels influence birthweight and 'catch-up' and 'catch-down' growth in a large contemporary cohort, 1207
- Stenhouse, E.A. *see* Janghorbani, M.
- Stenström, G. *see* Berger, B.
- Stephens, J.W. & Williams, R. Time to stop talking about screening for diabetes?, 1163
- Stephens, J.W. *see* Ashwell, S.G.
- Stephenson, C.R. *see* Feltbower, R.G.
- Stewart, J.A. *see* Gerstein, H.C.
- Stewart, M.W., Cirkel, D.T., Furuseth, K., Donaldson, J., Biswas, N., Starkie, M.G., Phenekos, C. & Hamann, A. Effect of metformin plus rosiglitazone compared with metformin alone on glycaemic control in well-controlled Type 2 diabetes, 1069
- Stock, S.A.K., Redaelli, M., Wendland, G., Civello, D. & Lauterbach, K.W. Diabetes—prevalence and cost of illness in Germany: a study evaluating data from the statutory health insurance in Germany, 299
- Stone, M.A. *see* Ali, S.
- Strassburger, K. *see* Hanifi-Moghaddam, P.
- Suárez, V. *see* Boronat, M.
- Su, D.-H., Lai, M.-Y. & Wu, H.-P. Liver failure in a patient receiving rosiglitazone therapy, 105
- Subramaniam, T. *see* Lim, S.C.
- Suehiro, A. *see* Satani, K.
- Sulli, N. & Shashaj, B. Long-term benefits of continuous subcutaneous insulin infusion in children with Type 1 diabetes: a 4-year follow-up, 900
- Sultan, A., Piot, C., Mariano-Goulart, D., Daures, J.P., Comte, F., Renard, E. & Avignon, A. Myocardial perfusion imaging and cardiac events in a cohort of asymptomatic patients with diabetes living in southern France, 410
- Sum, C.F. *see* Lim, S.C.
- Sun, G.Z. *see* Yang, J.K.
- Sun, L., Yang, Z., Jin, F., Zhu, X.Q., Qu, Y.C., Shi, X.H. & Wang, L. The Gly482Ser variant of the PPARGC1 gene is associated with Type 2 diabetes mellitus in northern Chinese, especially men, 1085
- Sun, T.B., Yang, C.C.H. & Kuo, T.B.J. Effect of hyperbaric oxygen on cardiac neural regulation in diabetic individuals with foot complications, 360
- Sundkvist, G. *see* Berger, B.
- Sundquist, K. & Li, X. Type 1 diabetes as a risk factor for stroke in men and women aged 15–49: a nationwide study from Sweden, 1261

- Sundvall, J. *see* Borodulin, K.
 Suraci, C. *see* Pitocco, D.
 Sutherland, M.G. *see* Gill, E.A.
 Sutton, S. *see* Farmer, A.
 Szczawinska, K. *see* Dworacka, M.
- Takada, H. *see* Tsutsumi, Y.
 Takata, K. *see* Hiromatsu, Y.
 Taketa, K. *see* Miyamura, N.
 Takiyama, Y. *see* Muroi, R.
 Tan, H.H. *see* Lim, S.C.
 Tan, I.K. *see* Lim, S.C.
 Tandon, N. *see* Vikram, N.K.
 Targher, G. *see* Molon, G.
 Targher, G., Bertolini, L., Padovani, R., Poli, F., Scala, L., Tessari, R., Zenari, L. & Falezza, G. Increased prevalence of cardiovascular disease in Type 2 diabetic patients with non-alcoholic fatty liver disease, 403
 Targher, G., Bertolini, L., Tessari, R., Zenari, L. & Arcaro, G. The International Diabetes Federation definition of the metabolic syndrome independently predicts future cardiovascular events in Type 2 diabetic patients. The Valpolicella Heart Diabetes Study, 1270
 Targher, G. Obesity and Diabetes, 1388
 Tarnow, L. *see* Lajer, M.
 Tauveron, I., Delcourt, I., Desbiez, F., Somda, F. & Thiéblot, P. Canine detection of hypoglycaemic episodes whilst driving, 335
 Taylor, H.R. *see* Phiri, R.
 Taylor, R. *see* Arun, C.S.
 Tee, W.W. *see* Lim, S.C.
 Temple, R.C. *see* Sampson, M.J.
 Ter Maaten, J.C. *see* Hagedoorn, M.
 Tessari, R. *see* Targher, G.
 The DIAMOND Project Group. Incidence and trends of childhood Type 1 diabetes worldwide 1990–1999, 857
 The ESRD Incidence Study Group. Divergent trends in the incidence of end-stage renal disease due to Type 1 and Type 2 diabetes in Europe, Canada and Australia during 1998–2002, 1364
 Thebault, L. *see* Schiele, F.
 Thiéblot, P. *see* Tauveron, I.
 Thomas, M.C., Tsalmandris, C., MacIsaac, R. & Jerums, G. Functional erythropoietin deficiency in patients with Type 2 diabetes and anaemia, 502
 Thomas, P. *see* Coppini, D.V.
 Thompson, C.H. *see* Johnson, N.A.
 Thompson, M.W. *see* Johnson, N.A.
 Thomsen, R.W. *see* Kruse, A.
 Thorsteinsson, B. *see* Akram, K.
 Todd, S. *see* Northam, E.A.
 Toh, V., Duncan, E., Lewis, N., Fichter, L. & Matthews, D.R. Ergotamine use in severe diabetic autonomic neuropathy, 574
 Tokunaga, H. *see* Miyamura, N.
 Tolis, G. *see* Angelopoulos, N.G.
 Tomelini, M. *see* Trento, M.
 Torlone, E. *see* Hoogma, R.P.L.M.
 Toth, E.L. *see* Johnson, J.A.
 Tovar, A. *see* Laffel, L.M.B.
 Tracy, R.P. *see* Davidson, M.
 Trautmann, M.E. *see* Linnebjerg, H.
- Trento, M., Tomelini, M., Lattanzio, R., Brancato, R., Coggiola, A., Benecchi, R., Scoccianti, L., Insacco, C., Bandello, F., Montanaro, M., Cavallo, F. & Porta, M. Perception of, and anxiety levels induced by, laser treatment in patients with sight-threatening diabetic retinopathy. A multicentre study, 1106
 Tsalmandris, C. *see* Thomas, M.C.
 Tsuchiya, T. *see* Shimizu, H.
 Tsutsumi, Y., Jie, X., Ihara, K., Nomura, A., Kanemitsu, S., Takada, H. & Hara, T. Phenotypic and genetic analyses of T-cell-mediated immunoregulation in patients with Type 1 diabetes, 1145
 Tuomilehto, J. *see* Borodulin, K.
 Tuomilehto, J. *see* Mager, U.
 Turner, C. *see* Wotherspoon, F.
 Turunen, J.A., Wessman, M., Kilpikari, R., Parkkonen, M., Forsblom, C. & Groop, P-H., on behalf of the FinnDiane Study Group. The functional variant -169C/T in the FCRL3 gene does not increase susceptibility to Type 1 diabetes, 925
 Tusek, C. *see* Bott, S.
 Tusié-Luna, M.T. *see* Navalón-García, J.
 Tuzcu, A.K. *see* Ozmen, S.
 Twomey, A.M. *see* Pegge, N.C.
- Umehara, A. *see* Satani, K.
 Unal, S. *see* Ertugrul, M.B.
 Uusitupa, M. *see* Mager, U.
- Vague, P. *see* Sejl, S.
 Vaidya, B. *see* Castleden, H.A.J.
 Valentine, W.J. *see* Roze, S.
 Valle, T.T. *see* Mager, U.
 van de Laar, F.A., van de Lisdonk, E.H., Lucassen, P.L.B.J., Stafleu, A., Mulder, J., van den Hoogen, H.J.M., Rutten, G.E.H.M. & van Weel, C. Eating behaviour and adherence to diet in patients with Type 2 diabetes mellitus, 788
 van de Lisdonk, E.H. *see* van de Laar, F.A.
 van den Hoogen, H.J.M. *see* van de Laar, F.A.
 van der Kallen, C.J.H. *see* Corpeleijn, E.
 van Esch, S.C.M., Cornel, M.C. & Snoek, F.J. Type 2 diabetes and inheritance: what information do diabetes organizations provide on the Internet?, 1233
 Van Le, K. *see* Norris, S.L.
 van Weel, C. *see* van de Laar, F.A.
 Varillas, V.F. *see* Boronat, M.
 Vaughton, K. *see* Pegge, N.C.
 Venugopal, H. *see* Baskar, V.
 Vering, T. *see* Schaller, H.C.
 Verza, M. *see* Marfella, R.
 Viberti, G., Lachin, J., Holman, R., Zinman, B., Haffner, S., Kravitz, B., Heise, M.A., Jones, N.P., O'Neill, M.C., Freed, M.I., Kahn, S.E. & Herman, W.H. for the ADOPT Study Group. A Diabetes Outcome Progression Trial (ADOPT): baseline characteristics of Type 2 diabetic patients in North America and Europe, 1289
 Vidya, M.-A. *see* Agius, E.
 Viggers, L. *see* Rosenfalck, A.M.
 Vignini, A. *see* Ferretti, G.
 Vikram, N.K., Tandon, N., Misra, A., Srivastava, M.C., Pandey, R.M., Mithal, A., Sharma, S., Ajmani, A., Madhu, S.V., Batra, C.M. & Gupta, N. Correlates of Type 2 diabetes mellitus in

- children, adolescents and young adults in north India: a multisite collaborative case-control study, 293
- Villarreal-Molina, T. *see* Navalón-García, K.
- Visalli, N. *see* Pitocco, D.
- von Sengbusch, S., Müller-Godeffroy, E., Häger, S., Reintjes, R., Hiort, O. & Wagner, V. Mobile diabetes education and care: intervention for children and young people with Type 1 diabetes in rural areas of northern Germany, 122
- Vuillermin, P.J. *see* Ismail, D.
- Wach, P. *see* Schaller, H.C.
- Wagenknecht, L.E. *see* Bowden, D.W.
- Wagenknecht, L.E. *see* Burdon, K.P.
- Wagenknecht, L.E. *see* Pradhan, A.D.
- Wagner, V. *see* von Sengbusch, S.
- Walden, E. *see* Sampson, M.J.
- Walker, B.R. Cortisol—cause and cure for metabolic syndrome?, 1281
- Walker, M. *see* Castleden, H.A.J.
- Wallén, N.H. *see* Yngen, M.
- Waller, A. *see* Franklin, V.L.
- Wallymahmed, M. *see* Woodward, A.
- Walters, M.R. *see* Nazir, F.S.
- Wang, C. *see* Hu, C.
- Wang, H. *see* Liu, L.
- Wang, L. *see* Sun, L.
- Wang, L. *see* Yang, J.K.
- Ware, R. *see* Xu, F.
- Watanabe, G. *see* Watarai, A.
- Watarai, A., Nakashima, E., Hamada, Y., Watanabe, G., Naruse, K., Miwa, K., Kobayashi, Y., Kamiya, H., Nakae, M., Hamajima, N., Sekido, Y., Niwa, T., Oiso, Y. & Nakamura, J. Aldose reductase gene is associated with diabetic macroangiopathy in Japanese Type 2 diabetic patients, 894
- Webb, D. *see* Ferriss, J.B.
- Weber, M.M. *see* Pfützner, J.
- Wegner, L., Andersen, G., Albrechtsen, A., Sparso, T., Glümer, C., Borch-Johnsen, K., Jørgensen, T., Hansen, T. & Pedersen, O. Large-scale study of the -232C > G polymorphism of *PCK1* in Type 2 diabetes, 1140
- Weidner, G. *see* Pradhan, A.D.
- Weill, A. *see* Cogneau, J.
- Wender-Ozegowska, E. *see* Dworacka, M.
- Wendland, G. *see* Stock, S.A.K.
- Weng, J. *see* Zhang, D.
- Wentzell, K. *see* Laffel, L.M.B.
- Wessman, M. *see* Bahr, C.
- Wessman, M. *see* Turunen, J.A.
- Whincup, P., Gilg, J., Owen, C., Cook, D. & Alberti, K.G.M.M. Authors' reply, 336
- Whincup, P.H. *see* Shultis, W.A.
- White, K.E. Research into the glomerular podocyte—is it relevant to diabetic nephropathy?, 715
- Whitehead, M. *see* Milton, B.
- Wiefels, K.J. *see* Hoogma, R.P.L.M.
- Wild, S. *see* Forouhi, N.
- Wild, S.H. *see* Forouhi, N.G.
- Wilding, J. *see* Woodward, A.
- Wilinska, M.E. *see* Schaller, H.C.
- Williams, A.J.K. *see* Castleden, H.A.J.
- Williams, J.S. *see* Raji, A.
- Williams, K. *see* Daly, M.E.
- Williams, R. *see* Deakin, T.A.
- Williams, R. *see* Stephens, J.W.
- Wilson, A.W. *see* Franklin, V.L.
- Winiarska, H. *see* Dworacka, M.
- Wise, S., Chien, J., Yeo, K. & Richardson, C. Smoking enhances absorption of insulin but reduces glucodynamic effects in individuals using the Lilly-Dura inhaled insulin system, 510
- Wolf, O.T. *see* Lobnig, B.M.
- Wolford, J.K., Yeatts, K.A., Eagle, A.R.R., Nelson, R.G., Knowler, W.C. & Hanson, R.L. Variants in the gene encoding aldose reductase (*AKR1B1*) and diabetic nephropathy in American Indians, 367
- Woodward, A. *see* Daousi, C.
- Woodward, A., Wallymahmed, M., Wilding, J. & Gill, G. Successful cardiovascular risk reduction in Type 2 diabetes by nurse-led care using an open clinical algorithm, 780
- Wootton, S.A. *see* Masding, M.G.
- Wotherspoon, F., Laight, D.W., Browne, D.L., Turner, C., Meeking, D.R., Allard, S.E., Munday, L.J., Shaw, K.M. & Cummings, M.H. Plasma homocysteine, oxidative stress and endothelial function in patients with Type 1 diabetes mellitus and microalbuminuria, 1350
- Wright, D.E. *see* Stenhouse, E.
- Wright, J. *see* Ismail, H.
- Wu, B.Y. *see* Yang, J.K.
- Wu, H.-P. *see* Su, D.-H.
- Wu, Y.S. *see* Lim, S.C.
- Xiang, K. *see* Hu, C.
- Xiu, L. *see* Zhang, D.
- Xu, F., Yin, X.M., Zhang, M., Leslie, E., Ware, R. & Owen, N. Family average income and diagnosed Type 2 diabetes in urban and rural residents in regional mainland China, 1239
- Xu, X. *see* Yang, J.K.
- Xu, X.H. *see* Lim, S.C.
- Xu, X.P. *see* Yang, J.K.
- Yagyu, H. *see* Muroi, R.
- Yale, J.-F. *see* Gerstein, H.C.
- Yamada, K. *see* Hiromatsu, Y.
- Yamasaki, Y. *see* Nakatani, Y.
- Yang, C.C.H. *see* Sun, T.B.
- Yang, G.R. *see* Yang, J.K.
- Yang, J.K., Feng, Y., Yuan, M.Y., Yuan, S.Y., Fu, H.J., Wu, B.Y., Sun, G.Z., Yang, G.R., Zhang, X.L., Wang, L., Xu, X., Xu, X.P. & Chan, J.C.N. Plasma glucose levels and diabetes are independent predictors for mortality and morbidity in patients with SARS, 623
- Yang, Z. *see* Sun, L.
- Yap, E. *see* Lim, S.C.
- Yeatts, K.A. *see* Wolford, J.K.
- Yeo, K. *see* Wise, S.
- Yin, X.M. *see* Xu, F.
- Yngen, M., Östenson, C.-G., Hjemdahl, P. & Wallén, N.H. Meal-induced platelet activation in Type 2 diabetes mellitus: effects of treatment with repaglinide and glibenclamide, 134
- Yoon, T.H. *see* Ahn, J.H.
- Yoshiuchi, K. *see* Nakatani, Y.

- Young, B. & Dean, J.D. Guest editorial, 1277
Young, D. *see* Arun, C.S.
Yu, Q. *see* Liu, L.
Yu, Y. *see* Lyons, T.J.
Yuan, M.Y. *see* Yang, J.K.
Yuan, S.Y. *see* Yang, J.K.
- Zaki, M.Y. *see* Mojiminiyi, O.A.
Zakrzewska, K. *see* Roze, S.
Zawiejska, A. *see* Dworacka, M.
Zenari, L. *see* Molon, G.
Zenari, L. *see* Targher, G.
Zervas, A. *see* Angelopoulos, N.G.
Zhang, C. *see* Zhang, D.
Zhang, D., Zhou, Z., Li, L., Weng, J., Huang, G., Jing, P., Zhang, C., Peng, J. & Xiu, L. Islet autoimmunity and genetic mutations in Chinese subjects initially thought to have Type 1B diabetes, 67
Zhang, H.-L. *see* Jian, W.-X.
Zhang, M. *see* Hughes, R.C.E.
Zhang, M. *see* Xu, F.
Zhang, R. *see* Hu, C.
Zhang, S.X. *see* Liu, L.
Zhang, X. *see* Norris, S.L.
Zhang, X.L. *see* Yang, J.K.
Zhao, Y. *see* Jian, W.-X.
Zheng, D. *see* Lyons, T.J.
Zheng, S. *see* Jian, W.-X.
Zhou, Z. *see* Zhang, D.
Zhu, J. *see* Davidson, M.
Zhu, X.Q. *see* Sun, L.
Ziemer, D.C. *see* Rhee, M.K.
Zimmet, P. *see* Alberti, K.G.M.M.
Zinman, B. *see* Shah, B.R.
Zinman, B. *see* Viberti, G.

Subject Index

- 232C/ G polymorphism, 419
- 1, 171
- 11 β -hydroxysteroid dehydrogenase, 1281
- 5-anhydro-D-glucitol, 171
- α -glucosidase inhibitor, 253
- abdominal obesity, 1025
- accumulation, 522
- aceruloplasminaemia, 1136
- acute hyperglycaemia, 171
- acute infectious conjunctivitis, 393
- adherence, 40, 265, 720, 795
- adiponectin, 253, 436, 660
- adolescence, 830, 1225, 1332
- adolescents, 293, 900, 1155, 1222
- ADOPT, 1289
- adult diabetes, 1186, 1247
- adults, 445, 1165
- African-Americans, 698
- ageing, 1174
- aids, 1110
- albuminuria, 185, 502
- alcohol, 795, 830
- alcohol drinking, 690
- aldose reductase, 367, 894
- aldosterone synthase, 675
- algorithms, 90
- Anaemia, 839
- angiotensin, 53
- antihypertensives, 426
- antimicrobials, 348
- antioxidants, 258
- anxiety, 917, 1106, 1110, 1381
- aortic stiffness, 1201
- arrhythmia, 207
- arteries, 873
- artificial β -cell, 90
- artificial pancreas, 1
- Asian diabetes, 1257
- Asian Indians, 293, 537, 666, 845
- association, 967
- association study, 228, 1140
- asymptomatic bacteriuria, 185
- atherosclerosis, 228, 609, 808, 894, 955
- ATP-sensitive potassium channel, 327
- attention, 26, 32
- automated insulin delivery, 1
- autonomic neuropathy, 360, 574, 873
- Avandamet, 1069
- β -hydroxybutyrate, 278
- β -thalassaemia major, 1327
- bacteriological analysis, 99
- basal insulin, 814
- behavioural, 485
- beliefs, 265
- binge eating, 917
- biomarkers, 1344
- birth weight, 1301
- birthweight, 1207
- blind, 1110
- blood glucose, 795, 1079
- blood pressure, 94, 557, 780
- BMI, 72
- body composition, 384, 1357
- body weight, 1079
- caffeine, 574
- calcitriol, 920
- cancer, 1186
- candidal carriage, 455
- candidate gene, 367
- cardiac events, 410
- cardiovascular disease, 1281
- cardiovascular disease, 403, 1186
- cardiovascular risk, 887
- cardiovascular risk factors, 537
- cardiovascular risk reduction, 780, 1117
- care delivery, 377
- carotid artery intima-media thickness, 609
- carotid atherosclerosis, 617
- carotid IMT, 845
- carotid stenosis, 1377
- case-control study, 393
- CD14, 72
- central obesity, 469
- cerebellar ataxia, 1136
- cerebral blood flow, 398
- cerebral oedema, 571
- CGMS, 1124
- chemokines, 156
- childhood and adolescence, 1216
- children, 86, 122, 293, 900, 1357
- children and young people, 935
- China, 1239
- Chinese, 67
- chronic pain, 1021
- chronic wounds, 594
- citations, 176
- clamp, 246
- clinical algorithm, 780
- clinical studies, 360
- coenzyme Q10, 1344
- cognitive function, 26
- cohort studies, 690
- colour tissue Doppler, 1201
- community health workers, 544
- co-morbid depression, 445
- co-morbidity, 1165
- compartmental models, 21
- congestive cardiac failure, 1313
- continuous glucose sensing, 1
- continuous subcutaneous infusion of insulin, 204
- continuous subcutaneous insulin infusion, 90, 141

- control algorithm, 1
- controlled trials, 1050
- coping, 1110
- coronary artery calcification, 1192
- coronary stenosis, 410
- cost, 299, 1100
- cost-effectiveness analysis, 164
- co-stimulatory molecule, 1145
- C-peptide, 1180
- C-reactive protein, 763, 1192
- CYP11B2, 675
- cytokines, 156
- death, 623
- decision aids, 912
- decision support technique, 912
- decision support tools, 21
- definition, 720
- delivery of health care, 164
- denaturing high-performance liquid chromatography, 681
- depression, 917, 1110, 1165, 1381
- diabetes, 21, 86, 128, 176, 189, 240, 377, 393, 410, 426, 436, 460, 485, 529, 574, 736, 763, 873, 982, 1008, 1106, 1130, 1239, 1252, 1344, 1370
- diabetes education, 271, 544
- diabetes mellitus, 148, 164, 171, 299, 323, 455, 623, 666, 955, 1117, 1339, 1377
- diabetes mellitus Type 1, 156
- diabetes mellitus Type 2, 419, 757
- diabetes risk factors, 803
- diabetes screening, 803
- diabetes self-management, 271
- diabetic complications, 185
- diabetic foot, 99, 360, 649
- diabetic foot ulcer, 341, 348
- diabetic ketoacidosis, 278, 571
- diabetic nephropathy, 331, 367, 675, 681, 1037
- diabetic neuropathy, 1016, 1021
- diabetic pregnancy, 223
- diabetic retinopathy, 60, 643, 867, 1029, 1106
- diagnosis, 341, 649, 982, 1257
- diagnostic criteria, 1319
- diet, 795, 1225
- dietary adherence, 788
- dietary carbohydrate, 15
- digital non-mydratic photography, 867
- digital retinal screening, 629
- drug utilization, 1117
- dual-energy X-ray absorptiometry scanning, 384
- duplex ultrasonography, 1377
- early onset diabetes, 1136
- eating behaviour, 788
- echocardiography, 1201
- economics, 1100
- education, 122, 935
- educational interventions, 485
- effectiveness, 1100
- efficacy, 246
- electrolytes, 1124
- endothelial dysfunction, 258, 398
- endothelium, 873, 1350
- end-stage renal disease, 1364
- epidemiological model, 189
- epidemiology, 148, 313, 857
- epidemiology prospective studies, 690
- erectile dysfunction, 198, 873
- ergotamine, 574
- ER Stress, 216
- erythropoietin, 839
- erythropoietin, 574
- ethnicity, 77, 94, 1233
- euglycaemic glucose clamp, 522
- exenatide, 240
- exercise, 795
- expectations, 1003
- extrapyramidal signs, 1136
- familial, 331
- familial inheritance, 834
- family average income, 1239
- family history, 1233
- fasting glucose, 1151, 1319
- fasting glycaemia, 1327
- fasting plasma glucose, 990
- fetal programming, 1213
- first hospital admission, 1261
- fludrocortisone, 574
- focus groups, 1222
- folate, 529
- follow-up study, 1261
- GAD antibody, 211
- gender, 907
- gene-environment interactions, 907
- gene mutation, 67
- general practice, 788, 803, 887, 996
- General Practice Research Database, 516
- genetic association, 1093
- genetic association studies, 419
- genetic polymorphism, 1307
- genetics, 228, 1037, 1140, 1257, 1295
- genetics of Type 2 diabetes, 685
- genotyping, 419
- geographical distribution, 857
- Germany, 299
- gestational, 460
- gestational diabetes, 306, 565, 1213, 1319
- gestational diabetes mellitus, 313
- ghrelin, 685
- glargine, 46, 743
- glibenclamide, 134, 327
- gliclazide, 246, 757
- glomerular filtration, 715
- glomerular filtration rate, 1057
- glomerulopathy, 715
- glucocorticoids, 1281
- glucodynamic, 510
- glucose, 1025
- glucose control, 90

- glucose dependent, 240
- glucose intolerance, 1174
- glucose metabolism, 1307
- glucose monitoring, 90
- glucose sensors, 21
- glucose tolerance, 306, 907, 1061, 1327
- glucose tolerance test, 982
- glutamic acid decarboxylase antibodies, 834
- glycaemic control, 207, 557, 830, 1050, 1069, 1216, 1225, 1247, 1381
- glycated haemoglobin, 944
- graft vs. host disease, 436
- Graves' disease, 211
- growth factors, 594
- guideline, 460, 579

- haemoglobin A_{1c}, 1289
- HbA_{1c}, 40, 285, 757, 900, 990, 1151, 1357
- HbA_{1c} levels, 1050
- HbA_{1c} targets, 1069
- health care delivery, 122
- health-care evaluation mechanisms, 912
- health insurance, 299
- health policy, 1247
- health research impact, 176
- heart-rate monitoring, 360
- hemoxygenase-1, 666
- hepatocyte growth factor, 617
- hepatocyte nuclear factor-1 α , 1295
- high fat diet/ insulin resistance, 1061
- hippocampus, 32
- HNF-1 β , 1301
- HOMA-IR index, 654
- homeostasis model assessment, 1327
- homocysteine, 529, 808, 1192, 1350
- hormone replacement therapy, 763
- hospital settings, 1106
- hospitalization, 377
- hyperglycaemia, 134
- hyperinsulinaemia, 77
- hypertension, 1130
- hypoglycaemia, 26, 141, 729, 750, 1155
- hypoglycaemia awareness, 750

- IGT, 845
- IL-18, 211, 235
- impaired fasting glucose, 148, 803
- impaired fasting glycaemia, 1130
- impaired glucose tolerance, 148, 565, 609, 1025, 1130
- IMT of the carotid artery, 228
- incidence, 857, 1364
- incretin, 240
- Indian and Pakistani, 1003
- infant growth, 1207
- infection, 341, 348
- inflammation, 763, 974
- inhaled insulin, 510
- inpatient bed occupancy, 1008
- insulin, 879, 1025
- insulin allergy, 204
- insulin analogue, 522
- insulin analogues, 285, 743, 879
- insulin aspart, 729
- insulin-dependent diabetes, 698
- insulin detemir, 522, 729
- insulin glargine, 46, 285, 736, 814, 879
- insulin-like growth factor-1, 814
- insulin-like growth factor binding protein 1, 814
- insulin pump, 1
- insulin pump therapy, 86, 900
- insulin resistance, 94, 223, 306, 469, 510, 537, 654, 1307, 1357
- insulin sensitivity, 384
- insulin therapy, 176, 750, 879, 1155, 1216
- insulin-treated adults, 271
- intelligence, 26
- intercellular adhesion molecule 1, 643, 1093
- intermittent claudication, 327
- International Diabetes Federation (IDF), 469
- Internet, 1233
- intervention, 113
- intramuscular triglyceride, 1061
- islet antibodies, 1180
- islet autoantibodies, 67

- ketosis, 1124
- kidney disease, 502
- Koreans, 198

- lactic acidosis, 432
- laser treatment, 60, 1106
- latent autoimmune diabetes in adults, 834
- lay health workers, 544
- leptin, 660
- Leu72Met, 685
- lifecourse, 821
- lifestyle, 720
- linkworker, 1003
- lipid metabolism, 907
- lipids, 15, 94, 557, 1029
- lipodystrophy, 436
- lipo-prostaglandin E₁, 1339
- lipoproteins, 768, 955
- lispro, 743
- low-density lipoproteins, 808
- low-fat diet, 384

- macroangiopathy, 894
- macrosomia, 460
- magnesium supplementation, 1050
- magnetic resonance imaging, 32
- management, 1252
- Maori, 1313
- maternal glucose, 1207
- matrix metalloproteinases, 594
- maturity-onset diabetes of the young 3 mutation, 1295
- mean plasma glucose, 1151
- measurement, 720
- medication compliance, 265
- memory, 32
- meta-analysis, 1050

- metabolic control, 113
metabolic factors, 1016
metabolic syndrome, 403, 441, 469, 845, 996, 1192, 1281
metformin, 128, 223, 318, 323, 432, 757, 1069, 1079
methods, 912
microalbuminuria, 426, 1057
microvascular complications, 367
microvascular disease, 360
microvolt T-wave alternans, 207
midodrine, 574
MODY, 1257
mortality, 516, 698, 1186, 1313
mortality rates, 698
MTHFR, 529
multicomponent, 485
multiple daily injections, 141
myocardial infarction, 974, 1313, 1370
myocardial perfusion imaging, 410
- NADH oxidase, 1037
NADPH oxidase, 666
nateglinide, 757
nationwide study, 1261
natural history, 1021
needle puncture, 99
neighbourhood deprivation, 313
neonatal diabetes, 1301
neonatal morbidity, 318
nephropathy, 502, 1057, 1093
new definition, 469
nicotinamide, 920
nitric oxide, 398, 594
nitric oxide synthase, 398
non-alcoholic fatty liver disease, 403
non-HDL-cholesterol, 1029
normal glucose tolerance, 887
northern Chinese, 1085
NPH insulin, 814
nuclear magnetic resonance, 955
nurse-led clinics, 780
- obesity, 15, 94, 128, 441, 565, 907
obstetrics, 481
octreotide, 574
oestrogens, 768
offspring, 565
oral agents, 40
oral candidosis, 455
oral glucose-lowering drugs, 736
oral glucose tolerance test, 654, 967
ORP150, 216
osteomyelitis, 99, 649
outcome, 1100
overprotection, 271
overweight, 1079
oxidative damage, 808
oxidative stress, 258, 666, 974, 1344, 1350
- paediatrics, 1257
pancreatic B-cell function, 1180
pancreatic development, 1301
pancreatic venous drainage, 654
parents, 1222
partners, 271
patient-centred care, 944
patient participation, 164
patient symptoms, 60
PD-1, 1145
peripheral vascular disease, 327
peroxisome proliferator-activated receptor γ co-activator-1, 1085
pharmacodynamics, 522
pharmacokinetic, 510
pharmacokinetics, 323
phenotype, 1289
phosphoenolpyruvate carboxykinase, 419
physical activity, 1025
pioglitazone, 246, 436, 537
plasminogen activator inhibitor type 1, 887
platelets, 134
polaroid non-mydratic photography, 867
polymorphism, 72, 211, 235, 635, 643, 675, 685, 894, 1085
portal venous drainage, 654
postchallenge glucose, 1174
postchallenge hyperglycemia, 982
postprandial, 240, 768
postprandial glucose, 743, 990, 1151
postprandial lipoproteins, 258
postural hypotension, 574
PPAR delta, 1307
PPARGC1B, 635
pre-diabetes, 156
preeclampsia, 318
pregnancy, 171, 306, 318, 323, 426, 460, 481, 1213
premature stroke, 1261
preterm, 426
prevalence, 148, 189, 198, 299, 441, 445, 982, 1165
primary care, 86, 516, 851, 1117
primary prevention, 1233
prognosis, 623
promoter, 235
proteinuria, 715
psychiatric morbidity, 113
psychological support, 917
psychology, 935
psychosocial, 485
pulmonary oedema, 571
- qualitative methods, 1222
quality assurance, 629
quality of care, 851
quality of life, 60, 141
QUICKI index, 654
- race, 1289
randomized controlled trial, 736, 944
red wine, 974
regulatory T cell, 1145
renal failure, 432
repaglinide, 134

- research evaluation, 176
- resistin, 660
- rest homes, 1252
- retinal pigment degeneration, 1136
- reversibility, 1016
- risk, 393
- risk factor, 313
- risk factors, 198, 698, 750, 1377
- risk score, 996, 1370
- rural, 122

- safety, 246
- SARS, 623
- screening, 867, 996, 1319
- self-efficacy, 1332
- self-management, 544
- self-monitor blood glucose, 1247
- sensitivity, 629
- sequence variants, 681
- serum, 156
- service users, 917
- services, 1003
- sex hormone-binding globulin, 306
- sick day management, 278
- silent myocardial ischaemia, 775, 1186
- simultaneous pancreas-kidney transplantation, 654
- single nucleotide polymorphism, 967, 1093
- single-strand conformation polymorphism, 643
- smoking, 510
- social class, 313
- social consequences, 821
- social support, 1332
- socio-economic status, 1239
- South Asians, 845
- South Indians, 1029
- specialist care, 1117
- specialist clinics, 851
- specialist nurse, 1008
- specificity, 629
- spinal cord infarction, 571
- standardized test meal, 743
- steroid, 623
- stress hyperglycaemia, 1370
- stroke, 529, 894
- structured group education, 944
- subclinical disease, 955
- sudden hearing loss, 1339
- sulphonylureas, 128
- systematic review, 341, 348, 609, 821
- systemic lupus erythematosus, 1145

- temporal trends, 1364
- text message, 1332
- thiazolidinedione, 253

- timing, 46
- tissue-type plasminogen activator, 887
- TNF- α , 660
- tocopherol, 1344
- total prevalence, 189
- treatment, 720, 1021
- trend, 857
- triglyceride, 72, 635
- triglyceride-rich lipoproteins, 258
- triglycerides, 1029
- tumour necrosis factor- α , 253
- Type 1 diabetes, 26, 32, 46, 53, 113, 122, 235, 278, 285, 384, 445, 481, 557, 675, 681, 698, 729, 808, 834, 857, 879, 920, 935, 1079, 1124, 1145, 1155, 1201, 1216, 1222, 1225, 1261, 1364, 1381
- Type 1 diabetes mellitus, 141, 571, 821, 1093, 1332, 1350
- Type 1 diabetic children, 660
- Type 1 DM, 216
- Type 1B diabetes mellitus, 67
- Type 2 diabetes, 15, 40, 185, 204, 207, 228, 246, 265, 293, 318, 398, 403, 432, 481, 502, 516, 579, 609, 750, 775, 788, 834, 845, 944, 990, 996, 1003, 1025, 1069, 1100, 1140, 1192, 1233, 1289, 1313, 1357, 1364
- Type 2 diabetes mellitus, 72, 134, 198, 253, 327, 617, 635, 690, 768, 839, 967, 1085, 1165
- Type 2 diabetes susceptibility, 1295

- ulcer swabbing, 99
- undiagnosed diabetes, 803
- university students, 1155
- urban Nepal, 1130
- urine ketones, 278
- utility, 1100

- variation in care, 377
- vascular complications, 1100
- vascular disease, 1350
- vascular dysfunction, 537
- vascular endothelium, 53
- vibration perception threshold, 1016
- visfatin, 967
- visual impairment, 60

- weight, 900
- weight gain, 557
- weight loss, 15
- within-person variation, 729
- women, 77
- women's health, 763
- wound healing, 594

- xylosyltransferase I, 681

- young adults, 293



